

AD-A199 809

## REPORT DOCUMENTATION PAGE

②

1a. SECURITY CLASSIFICATION AUTHORITY UNCLASSIFIED		1b. RESTRICTIVE MARKINGS							
2a. DECLASSIFICATION/DOWNGRADING SCHEDULE NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited							
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 238		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TX- 83 - 0965							
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM							
6c. ADDRESS (City, State and ZIP Code) Statistics Department CB #3260, Phillips Hall Chapel Hill, NC 27599-3260		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448							
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144							
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		10. SOURCE OF FUNDING NOS. <table border="1"><tr><td>PROGRAM ELEMENT NO. 6.1102F</td><td>PROJECT NO. 2304</td><td>TASK NO. ALC</td><td>WORK UNIT NO. Leave blank</td></tr></table>		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304	TASK NO. ALC	WORK UNIT NO. Leave blank		
PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304	TASK NO. ALC	WORK UNIT NO. Leave blank						
11. TITLE (Include Security Classification) A Langevin-type Stochastic Differential Equation on a Space of Generalized functionals									
12. PERSONAL AUTHOR(S) Kallianpur, G. and Mitoma, I.									
13a. TYPE OF REPORT preprint	13b. TIME COVERED FROM 9/1/87 TO 8/31/88	14. DATE OF REPORT (Yr., Mo., Day) 1988, August	15. PAGE COUNT 48						
16. SUPPLEMENTARY NOTATION									
COSATI CODES <table border="1"><tr><td>FIELD</td><td>GROUP</td><td>SUB. GR.</td></tr><tr><td>XXXX</td><td>XXXXXXXXXX</td><td></td></tr></table>		FIELD	GROUP	SUB. GR.	XXXX	XXXXXXXXXX		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Key words and phrases: Weak solution, SDE, Frechet derivative, generalized functional space, central limit theorem, system of neurons.	
FIELD	GROUP	SUB. GR.							
XXXX	XXXXXXXXXX								
19. ABSTRACT (Continue on reverse if necessary and identify by block number) <p>Abstract. Let <math>E'</math> be the dual of a nuclear Fréchet space <math>E</math> and <math>L^*(t)</math> the adjoint operator of <math>L(t)</math> which has a formal expression:</p> $L(t) = \sum_1 a_i(t,x) \frac{\partial^2}{\partial x_i^2} + b_i(t,x) \frac{\partial}{\partial x_i}.$ <p>It is shown that the weak solution of a stochastic differential equation</p> $dX(t) = dW(t) + L^*(t)X(t)dt.$									
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT CLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED							
22a. NAME OF RESPONSIBLE INDIVIDUAL Major Brian W. Woodruff		22b. TELEPHONE NUMBER (Include Area Code) (202) 767-5026	22c. OFFICE SYMBOL AFOSR/NM						

ORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGEDTIC  
ELECTE  
OCT 06 1988  
D  
R E


19. Abstract (cont'd).

(UNCLASSIFIED)

exists uniquely on a generalized functional space on  $E'$  which is an appropriate model for the central limit theorem for an interacting system of spatially extended neurons. Applications to the latter problem are discussed.

# CENTER FOR STOCHASTIC PROCESSES

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina



A LANGEVIN-TYPE STOCHASTIC DIFFERENTIAL EQUATION ON A  
SPACE OF GENERALIZED FUNCTIONALS

by

G. Kallianpur

and

I. Mitoma

Technical Report No. 238

August 1988

88 10 5 153

163. M. O'Sullivan and T.R. Fleming, Statistics for the two-sample survival analysis problem based on product limit estimators of the survival functions, Nov. 86.
164. F. Avram, On bilinear forms in Gaussian random variables, Toeplitz matrices and Parseval's relation, Nov. 86.
165. D.B.H. Cline, Joint stable attraction of two sums of products, Nov. 86. *J. Multivariate Anal.*, 25, 1988, 272-285.
166. R.J. Wilson, Model fields in crossing theory—a weak convergence perspective, Nov. 86.
167. D.B.H. Cline, Consistency for least squares regression estimators with infinite variance data, Dec. 86.
168. L.L. Campbell, Phase distribution in a digital frequency modulation receiver, Nov. 86.
169. B.C. Nguyen, Typical cluster size for 2-dim percolation processes, Dec. 86. *J. Statist. Physics*, to appear.
170. H. Oodaira, Freidlin-Wentzell type estimates for a class of self-similar processes represented by multiple Wiener integrals, Dec. 86.
171. J. Nolan, Local properties of index- $\beta$  stable fields, Dec. 86. *Ann. Probability*, to appear.
172. R. Menich and R.F. Serfozo, Optimality of shortest queue routing for dependent service stations, Dec. 86.
173. F. Avram and M.S. Taqqu, Probability bounds for M-Skorohod oscillations, Dec. 86.
174. F. Moricz and R.L. Taylor, Strong laws of large numbers for arrays of orthogonal random variables, Dec. 86.
175. G. Kallianpur and V. Perez-Abreu, Stochastic evolution equations driven by nuclear space valued martingales, Apr. 87. *Appl. Math. Optimization*, 17, 1988, 237-272.
176. E. Merzbach, Point processes in the plane, Feb. 87.
177. Y. Kasahara, M. Maejima and W. Vervaat, Log fractional stable processes, March 87.
178. G. Kallianpur, A.G. Mijne and H. Niemi, On the prediction theory of two parameter stationary random fields, March 87. *J. Multivariate Anal.*, 1988, to appear.
179. R. Brigola, Remark on the multiple Wiener integral, Mar. 87.
180. R. Brigola, Stochastic filtering solutions for ill-posed linear problems and their extension to measurable transformations, Mar. 87.
181. G. Samorodnitsky, Maxima of symmetric stable processes, Mar. 87.
182. H.L. Hurd, Representation of harmonizable periodically correlated processes and their covariance, Apr. 87.
183. H.L. Hurd, Nonparametric time series analysis for periodically correlated processes, Apr. 87.

184. T. Mori and H. Oodaira, Freidlin-Wentzell estimates and the law of the iterated logarithm for a class of stochastic processes related to symmetric statistics, May 87.
185. R.F. Serfozo, Point processes, May 87. *Operations Research Handbook on Stochastic Processes*, to appear.
186. Z.D. Bai, W.Q. Liang and W. Vervaat, Strong representation of weak convergence, June 87.
187. O. Kallenberg, Decoupling identities and predictable transformations in exchangeability, June, 87.
188. O. Kallenberg, An elementary approach to the Daniell-Kolmogorov theorem and some related results, June 87. *Math. Nachr.*, to appear.
189. G. Samorodnitsky, Extrema of skewed stable processes, June 87. *Stochastic Proc. Appl.*, to appear.
190. D. Nualart, M. Sanz and M. Zakai, On the relations between increasing functions associated with two-parameter continuous martingales, June 87.
191. F. Avram and M. Taqqu, Weak convergence of sums of moving averages in the  $\alpha$ -stable domain of attraction, June 87.
192. M.R. Leadbetter, Harald Cramér (1893-1985), July 87. *ISI Review*, 56, 1988, 89-97.
193. R. LePage, Predicting transforms of stable noise, July 87.
194. R. LePage and B.M. Schreiber, Strategies based on maximizing expected log, July 87.
195. J. Rosinski, Series representations of infinitely divisible random vectors and a generalized shot noise in Banach spaces, July 87.
196. J. Szulga, On hypercontractivity of  $\alpha$ -stable random variables,  $0 < \alpha < 2$ , July 87.
197. I. Kuznezova-Shoipo and S.T. Rachev, Explicit solutions of moment problems I, July 87. *Probability Math. Statist.*, 10, 1989, to appear.
198. T. Hsing, On the extreme order statistics for a stationary sequence, July 87.
199. T. Hsing, Characterization of certain point processes, Aug. 87. *Stochastic Proc. Appl.*, 26, 1987, 297-316.
200. J.P. Nolan, Continuity of symmetric stable processes, Aug. 87.
201. M. Marques and S. Cambanis, Admissible and singular translates of stable processes, Aug. 88.
202. O. Kallenberg, One-dimensional uniqueness and convergence criteria for exchangeable processes, Aug. 87. *Stochastic Proc. Appl.*, 28, 1988, 159-183.
203. R.J. Adler, S. Cambanis and G. Samorodnitsky, On stable Markov processes, Sept. 87.
204. G. Kallianpur and V. Perez-Abreu, Stochastic evolution equations driven by nuclear space valued martingales, Sept. 87. *Appl. Math. Optimization*, 17, 1988, 237-272.
205. R.L. Smith, Approximations in extreme value theory, Sept. 87.

A Langevin-type Stochastic Differential Equation on a  
Space of Generalized Functionals

Gopinath Kallianpur\*  
Department of Statistics  
University of North Carolina  
Chapel Hill, NC 27599-3260 USA

and

Itaru Mitoma\*  
Department of Mathematics  
Hokkaido University  
Sapporo 060 Japan



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

**Abstract.** Let  $E'$  be the dual of a nuclear Fréchet space  $E$  and  $L^*(t)$  the adjoint operator of  $L(t)$  which has a formal expression:

$$L(t) = \sum_i a_i(t, x) \frac{\partial^2}{\partial x_i^2} + b_i(t, x) \frac{\partial}{\partial x_i}.$$

It is shown that the weak solution of a stochastic differential equation:

$$dX(t) = dW(t) + L^*(t)X(t)dt,$$

exists uniquely on a generalized functional space on  $E'$  which is an appropriate model for the central limit theorem for an interacting system of spatially extended neurons. Applications to the latter problem are discussed.

**Key words and phrases:** Weak solution, SDE, Fréchet derivative, generalized functional space, central limit theorem, system of neurons.

\*Research partially supported by the Air Force Office of Scientific Research Contract No. F49620 85 C 0144.

## 51. Introduction

Recently, Deuschel [4] has obtained a fluctuation result for a system of lattice valued diffusion processes. The result obtained is similar to the ones for mean-field interacting particle diffusions treated in a number of papers, [2,3,8,9,16,23]. In another direction, Kallianpur and Wolpert [11] have introduced a class of stochastic differential equations (SDE's) governing nuclear space valued processes as a model for voltage potentials for spatially extended neurons. The present paper is motivated by both the above problems, especially, the problem of interacting systems of neurons. The techniques developed in this paper enable us to prove a general result which yields a central limit theorem for such systems. It also provides another approach to the fluctuation theorem in [4]. In addition, the identification problem of the limit measures leads us to discuss the uniqueness of weak solutions of the SDE, formally expressed by

$$dX(t) = dW(t) + L^*(t)X(t)dt.$$

A precise meaning to the above equation is given by equation (1.1) below.

Our aim is to find a suitable space  $\mathfrak{D}_E$ , of smooth functionals on the dual nuclear space  $E'$  and to solve the SDE on the dual space  $\mathfrak{D}_E'$ , which is appropriate for the central limit theorems we have in mind.

We will proceed to explain the setting: A stochastic process  $X_F(t)$  defined on a complete probability space  $(\Omega, \mathcal{F}, P)$  indexed by elements in  $\mathfrak{D}_E$ , is called an  $\mathcal{V}(\mathfrak{D}_E)$ -process if  $X_F(t)$  is a real stochastic process for any fixed  $F \in \mathfrak{D}_E$ , and  $X_{\alpha F + \beta G}(t) = \alpha X_F(t) + \beta X_G(t)$  almost surely for each real numbers  $\alpha, \beta$  and elements of  $F, G \in \mathfrak{D}_E$ , and further  $E[X_F(t)^2]$  is continuous with respect to  $F$  on  $\mathfrak{D}_E$ , [10].  $X_F(t)$  is called continuous if  $\lim_{t \rightarrow s} E[(X_F(t) - X_F(s))^2] = 0$  for each  $F \in \mathfrak{D}_E$ . Let  $W_F(t)$  be an  $\mathcal{V}(\mathfrak{D}_E)$ -Wiener process, i.e. such that for any fixed

$F \in \mathfrak{D}_E$ .  $W_F(t)$  is a real continuous Gaussian additive process with mean 0.

We will prove that a unique continuous  $\mathfrak{L}(\mathfrak{D}_E)$ -process solution  $X_F(t)$  exists for the following equation with given initial value  $X_F(0)$ :

$$(1.1) \quad dX_F(t) = dW_F(t) + X_{L(t)}F(t)dt.$$

Roughly speaking, if  $L(t)$  generates the strongly continuous Kolmogorov evolution operator  $U(t,s)$  from  $\mathfrak{D}_E$  into itself, the unique solution for (1.1) can be given as follows:

$$X_F(t) = X_{U(t,0)}F(0) + W_F(t) + \int_0^t W_{L(s)}U(t,s)F(s)ds.$$

We will now begin by giving the precise definitions of the operator  $L(t)$  and the space  $\mathfrak{D}_E$ . Let  $E$  be a nuclear Fréchet space whose topology is defined by an increasing sequence of Hilbertian semi-norms  $\|\cdot\|_1 \leq \|\cdot\|_2 \leq \dots \leq \|\cdot\|_p \leq \dots$ . As usual let  $E'$  be the dual space,  $E'_p$  the completion of  $E$  by the  $p$ -th semi-norm  $\|\cdot\|_p$  and  $E'_p$  the dual space of  $E_p$ . Then we have

$$E = \bigcap_{p=0}^{\infty} E_p \quad \text{and} \quad E' = \bigcup_{p=0}^{\infty} E'_p.$$

Let  $K$  be a separable Hilbert space with norm  $\|\cdot\|_K$  and  $F$  a mapping from  $E'$  into  $K$ . Then  $F$  is said to be  $E'_p$ -Fréchet differentiable if for every  $x \in E'$ , we have a bounded linear operator  $\mathfrak{D}_p F(x)$  from  $E'_p$  into  $K$  such that

$$\lim_{t \rightarrow 0} \frac{F(x+th) - F(x)}{t} = \mathfrak{D}_p F(x)(h), \quad h \in E'_p, \quad \text{in } K.$$

Suppose that  $F$  is  $E'_p$ -Fréchet differentiable for every integer  $p \geq 0$ . Then taking  $E' = \bigcup_{p=0}^{\infty} E'_p$  and the strong topology of  $E'$ , (which is equivalent to the inductive limit topology of  $E'_p$ ;  $p=0,1,2,\dots$ ), into account, we have a continuous linear operator  $DF(x)$  from  $E'$  equipped with the strong topology into

$K$  such that for any integer  $p \geq 0$ ,  $DF(x)(h) = \mathfrak{D}_p F(x)\{h\}$  for  $h \in E'_p$ . Hence, if  $F$  is  $n$ -times  $E'_p$ -Fréchet differentiable for every integer  $p \geq 0$ , we have a continuous  $n$ -linear operator  $D^n F(x)$  from  $\underbrace{E' \times E' \times \dots \times E'}_{n\text{-times}}$  into  $K$  such that the restriction of  $D^n F(x)$  on  $\underbrace{E'_p \times E'_p \times \dots \times E'_p}_{n\text{-times}} =$  the  $n$ -th  $E'_p$ -Fréchet derivative  $\mathfrak{D}_p^n F(x)$ . Then if  $F$  is infinitely many times  $E'_p$ -Fréchet differentiable for every integer  $p \geq 0$ , the Hilbert-Schmidt norm

$$\|D^n F(x)\|_{H.S.}^{(p)} = \left( \sum_{i_1, i_2, \dots, i_n=1}^{\infty} \|\mathfrak{D}_p^n F(x)(h_{i_1}^{(p)}, h_{i_2}^{(p)}, \dots, h_{i_n}^{(p)})\|_K^2 \right)^{1/2}$$

is finite for each integer  $n \geq 1$  and  $p \geq 0$ , where  $(h_j^{(p)})$  is a C.O.N.S., (complete orthonormal system), in  $E'_p$  [14].

From now on, we will often use the conventional notation such that  $\|D^0 F(x)\|_{H.S.}^{(p)} = \|F(x)\|_K$ .

Let  $\beta(t)$  be the standard  $E'$ -Wiener process such that for any  $\xi \in E$ ,  $\langle \beta(t), \xi \rangle$  is a 1-dimensional Brownian motion, with variance  $E[\langle \beta(t), \xi \rangle^2] = t \|\xi\|_0^2$ , where  $\langle x, \xi \rangle$ , ( $x \in E'$ ,  $\xi \in E$ ), denotes the canonical bilinear form on  $E' \times E$ .

Without loss of generality, we assume  $\beta(t)$  is an  $E'_1$ -valued Wiener process throughout this paper, [17], [18].

**Definition of  $L(t)$ .** For  $t > 0$  and  $x \in E'$ , let  $A(t, x)$  and  $B(t, \cdot)$  be continuous mappings from  $E'$  into itself such that the following conditions are satisfied.

(H1) There exists a natural number  $p_0$  such that  $A(t, x)$  maps  $E'_1$  into  $E'_{p_0}$ .  $B(t, \cdot)$  maps  $E'$  into  $E'_{p_0}$  and for each  $T > 0$ ,



$$\sup_{\substack{x \in E' \\ 0 \leq t \leq T}} \|A(t, x)\|_2^2 < \infty \quad \text{and} \quad \sup_{\substack{x \in E' \\ 0 \leq t \leq T}} \|B(t, x)\|_{-p_0} < \infty,$$

where  $\|\cdot\|_{-p}$  denotes the dual norm of  $E'_p$  and  $\|A(t, x)\|_2^2 = \sum_{j=1}^{\infty} \|A(t, x)h_j^{(0)}\|_{-p_0}^2$ .

(H2)  $A(t, x)$  and  $B(t, x)$  are infinitely many times  $E'_p$ -Fréchet differentiable for every integer  $p \geq 0$  such that for any  $T > 0$  and any integer  $n \geq 1$ ,

$$\sup_{\substack{x \in E' \\ 0 \leq t \leq T}} \|D^n(t, x)\|_{H.S.}^{(p)} < \infty \quad \text{and} \quad \sup_{\substack{x \in E' \\ 0 \leq t \leq T}} \|D^n B(t, x)\|_{H.S.}^{(p)} < \infty,$$

where  $\|D^n A(t, x)\|_{H.S.}^{(p)} = \left( \sum_{i_1, i_2, \dots, i_n=1}^{\infty} \|D^n A(t, x)(h_{i_1}^{(p)}, h_{i_2}^{(p)}, \dots, h_{i_n}^{(p)})\|_2^2 \right)^{1/2}$  and  $\|D^n B(t, x)\|_{H.S.}^{(p)} = \left( \sum_{i_1, i_2, \dots, i_n=1}^{\infty} \|D^n B(t, x)(h_{i_1}^{(p)}, h_{i_2}^{(p)}, \dots, h_{i_n}^{(p)})\|_{-p_0}^2 \right)^{1/2}$ .

(H3) For any integer  $n \geq 0$  and any  $T > 0$ , there exist  $\lambda(n, p, T) > 0$  and  $\lambda_1(n, p, T) > 0$  such that

$$\sup_{\substack{x \in E' \\ 0 \leq k \leq n}} \max \{ \|D^k A(t, x) - D^k A(s, x)\|_{H.S.}^{(p)}, \|D^k B(t, x) - D^k B(s, x)\|_{H.S.}^p \} \\ \leq \lambda_1(n, p, T) |t-s|^{\lambda(n, p, T)}, \quad 0 \leq s, t \leq T.$$

Then for any twice  $E'_p$ -Fréchet differentiable real valued functional  $F$  on  $E'$  for every  $p \geq 0$ , we put

$$(L(t)F)(x) = \frac{1}{2} \text{trace}_{E_0} D^2 F(x) \circ [A(t, x) \times A(t, x)] + DF(x)(B(t, x)),$$

where  $\text{trace}_{E_0} D^2 F(x) \circ [A(t, x) \times A(t, x)] = \sum_{j=1}^{\infty} D^2 F(x)(A(t, x)h_j^{(0)}, A(t, x)h_j^{(0)})$ .

**Definition of  $\mathfrak{L}_E$ .** For a real valued infinitely many times  $E'_p$ -Fréchet differentiable functional  $F$  on  $E'$  for every integer  $p \geq 0$ , we define the following semi-norms:

$$\|F\|_{p,q,n} = \sum_{k=0}^n \|F\|_{p,k}^{(q)},$$

where  $p \geq 0$ ,  $q \geq 0$  and  $n \geq 0$  are integers and

$$\|F\|_{p,k}^{(q)} = \sup_{x \in E'_p} e^{-\|x\|} {}^p D^k F(x) \|_{H.S.}^{(q)}.$$

For any natural number  $n$ , define

$$S(\mathbb{R}^n) = \{\phi(x) = h(x)\varphi(x); \varphi \in \mathcal{S}(\mathbb{R}^n)\},$$

where  $\varphi(x)$  is an element of the Schwartz space  $\mathcal{S}(\mathbb{R}^n)$  of rapidly decreasing  $C^\infty$ -functions on the  $n$ -dimensional Euclidean space  $\mathbb{R}^n$  and  $h(x)$ ,

$x=(x_1, x_2, \dots, x_n)$ , is a weight function such that  $h(x) = 1/g(x)$ .

$g(x) = \prod_{i=1}^n g_0(x_i)$ ,  $g_0(x_i) = \exp(-\sqrt{\int_{\mathbb{R}} |y| \rho(x_i - y) dy})$  and  $\rho(x)$  is the Friedrichs mollifier whose support is contained in  $[-1, 1]$ . Let  $\{\xi_j; j=1, 2, \dots\}$  be a countable dense subset of  $E$ . Define

$$C_{0,n}^\infty(E') = \{\phi(x) = \phi(\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_n \rangle); \phi \in S(\mathbb{R}^n)\}$$

and introduce the nuclear Fréchet topology on this space by the countably many semi-norms;

$$\|\phi\|_p = \sup_{\substack{x \in \mathbb{R}^n \\ 0 \leq k \leq p}} (1+|x|^2)^p |(\frac{d}{dx})^k (g(x)\phi(x))|, \quad p=0, 1, 2, \dots,$$

where  $(\frac{d}{dx})^k = \sum_{k_1+k_2+\dots+k_n=k} \frac{\partial^k}{\partial x_1^{k_1} \partial x_2^{k_2} \dots \partial x_n^{k_n}}$ . Then we have a fundamental space

$C_0^\infty(E') = \bigcup_{n=1}^\infty C_{0,n}^\infty(E')$  which is the strict inductive limit of nuclear Fréchet spaces  $C_{0,n}^\infty(E')$ .

For any integers  $p \geq 0$ ,  $q \geq 0$  and  $n \geq 0$ , let  $\mathfrak{D}_{p,q,n}$  be the completion of  $C_0^\infty(E')$  by the semi-norm  $\|\cdot\|_{p,q,n}$ . We define  $\mathfrak{D}_E = \bigcap_{p,q,n} \mathfrak{D}_{p,q,n}$  and introduce a

topology on  $\mathfrak{D}_E$ , by the countably many semi-norms  $\|\cdot\|_{p,q,n}$ ,  $p \geq 0$ ,  $q \geq 0$  and  $n \geq 0$ .

Then  $\mathfrak{D}_E$  becomes a complete separable metric space [6].

**Remark 1.** The definition of  $\mathfrak{D}_E$  is independent of the way of choosing a countable dense subset of  $E$ . We call a real valued functional expressed as  $\phi(x) = \phi(\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_n \rangle)$  by using some natural number  $n$ ,  $\xi_i \in E$ ,  $i=1,2,\dots,n$ , and  $\phi \in S(\mathbb{R}^n)$  a weighted Schwartz functional. Let  $\mathcal{S}$  be the set of all weighted Schwartz functionals,  $\mathcal{S}_{p,q,n}$  the completion of  $\mathcal{S}$  by  $\|\cdot\|_{p,q,n}$  and  $\mathfrak{D} = \bigcap_{p,q,n} \mathcal{S}_{p,q,n}$ , where  $p \geq 0$ ,  $q \geq 0$  and  $n \geq 0$  are integers. Then

$$\mathfrak{D} = \mathfrak{D}_E.$$

**Proof.** It is enough to show that  $\phi(x) = \phi(\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_m \rangle)$ ,  $\xi_i \in E$ ,  $\phi \in S(\mathbb{R}^m) \in \mathfrak{D}_{p,q,n}$ . By the nuclearity of  $E$ , we have a natural number  $r > \max\{p,q\}$  such that

$$(1.2) \quad \sum_{j=1}^{\infty} \|h_j^{(q)}\|_{-r}^2 < \infty$$

and since  $\{\xi_j\}$  is dense in  $E$ , for each  $i$ , there exists a sequence  $\{\xi_{i,k}\}$ ,  $\xi_{i,k} \in \{\xi_j\}$  such that

$$(1.3) \quad \lim_{k \rightarrow \infty} \|\xi_i - \xi_{i,k}\|_r = 0.$$

On the other hand,  $D^n \phi(x)(h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_n}^{(q)})$  is a finite sum of terms;

$$(1.4) \quad \frac{\partial^n}{\partial x_1^{n_1} \partial x_2^{n_2} \dots \partial x_m^{n_m}} \phi(\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_m \rangle) \langle h_{j_1}^{(q)} \rangle_{j_1} \cdot \xi_1 \rangle \\ \times \langle h_{j_2}^{(q)} \rangle_{j_2} \cdot \xi_1 \rangle \dots \langle h_{j_{n_1}}^{(q)} \rangle_{j_{n_1}} \cdot \xi_1 \rangle \langle h_{j_1}^{(q)} \rangle_{j_1} \cdot \xi_2 \rangle \langle h_{j_2}^{(q)} \rangle_{j_2} \cdot \xi_2 \rangle \dots \langle h_{j_{n_2}}^{(q)} \rangle_{j_{n_2}} \cdot \xi_2 \rangle$$

$$\dots \langle h_{j_1^{(m)}}^{(q)}, \xi_m \rangle \langle h_{j_2^{(m)}}^{(q)}, \xi_m \rangle \dots \langle h_{j_n^{(m)}}^{(q)}, \xi_m \rangle,$$

where  $n_1 + n_2 + \dots + n_m = n$ . Since

$$(1.5) \quad \left| \frac{\partial^\mu}{\partial x_1^{\mu_1} \partial x_2^{\mu_2} \dots \partial x_m^{\mu_m}} h(x) \right| \leq C_1 \exp \left( \sum_{i=1}^m \sqrt{|x_i|} \right).$$

Noticing  $\phi(x) = h(x)\varphi(x)$ ,  $\varphi \in \mathcal{V}(\mathbb{R}^m)$ , we have

$$(1.6) \quad \sup_k \sup_{x \in E'_p} e^{-\|x\|_p} \max \left\{ \left| \frac{\partial^n}{\partial x_1^{n_1} \partial x_2^{n_2} \dots \partial x_m^{n_m}} \phi(\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_m \rangle) \right| \right.$$

$$\left. \left| \frac{\partial^n}{\partial x_1^{n_1} \partial x_2^{n_2} \dots \partial x_m^{n_m}} \phi(\langle x, \xi_{1,k}^0 \rangle, \langle x, \xi_{2,k}^0 \rangle, \dots, \langle x, \xi_{m,k}^0 \rangle) \right| \right.$$

$$\left. \left| \frac{\partial^{n_1} \dots \partial^{n_{i-1}} \partial^{n_i+1} \partial^{n_{i+1}} \dots \partial^{n_m}}{\partial x_1^{n_1} \dots \partial x_{i-1}^{n_{i-1}} \partial x_i^{n_i+1} \partial x_{i+1}^{n_{i+1}} \dots \partial x_m^{n_m}} \phi(\langle x, \xi_{1,k}^0 \rangle, \langle x, \xi_{2,k}^0 \rangle, \dots \right. \right.$$

$$\left. \dots, \langle x, \xi_{i,k}^0 \rangle + \tau(\langle x, \xi_i \rangle - \langle x, \xi_{i,k}^0 \rangle), \langle x, \xi_{i+1} \rangle, \dots, \langle x, \xi_m \rangle) \right|$$

$$\times \|x\|_p^{-p}, \quad 0 \leq \tau \leq 1, \quad i=1, 2, \dots, m \} \leq C_2.$$

Setting  $\psi^{(k)}(x) = \phi(\langle x, \xi_{1,k}^0 \rangle, \langle x, \xi_{2,k}^0 \rangle, \dots, \langle x, \xi_{m,k}^0 \rangle)$  and using (1.2) - (1.6), we have

$$\lim_{k \rightarrow \infty} \|\phi - \psi^{(k)}\|_{p,q,n} = 0,$$

which completes the proof.

Here and in the sequel, we denote positive constants by  $C_1$  or, if necessary, by  $C_1(\tau_1, \tau_2, \dots)$ ,  $i=1, 2, \dots$ , in case they depend on the parameters  $\tau_1, \tau_2, \dots$ .

Before proceeding to the discussion of equation (1.1), the following remarks on the  $\mathcal{L}(\mathfrak{H}_E)$ -Wiener process are in order. Taking the continuity of  $W_F(t)$  and  $E[W_F(t)^2]$  with respect to the parameters  $t$  and  $F$  into account, we note that  $\sup_{0 \leq t \leq T} E[W_F(t)^2] < \infty$  and  $\sup_{0 \leq t \leq T} E[W_F(t)^2]$  is lower semi-continuous on  $\mathfrak{H}_E$ . Since  $\mathfrak{H}_E$  is a complete metric space, by the Banach-Steinhaus theorem we have some positive integers  $p_1, q_1$  and  $m_1$  such that

$$(1.7) \quad \sup_{0 \leq t \leq T} E[W_F(t)^2] \leq C_3(T) \|F\|_{p_1, q_1, m_1}^2.$$

Now given a functional  $V_t(F)$  such that for each  $t$  it is a positive definite quadratic form on  $\mathfrak{H}_E \times \mathfrak{H}_E$ , increasing and continuous in  $t$  and  $\sup_{0 \leq t \leq T} V_t(F) \leq C_4(T) \|F\|_{p, q, n}^2$  for some natural numbers  $p, q$  and  $n$ , we can construct a  $\mathfrak{H}_E$ -indexed Gaussian mean-zero continuous process  $W_F(t)$  with independent increments and variance  $V_t(F)$  by the Kolmogorov theorem, since  $V_{t \wedge s}(F)$  is positive definite quadratic form with respect to  $(t, F)$ ,  $t \in [0, \infty)$ ,  $F \in \mathfrak{H}_E$ . Here  $t \wedge s = \min \{t, s\}$ .

## §2. Existence and Uniqueness of solutions of the SDE

Let  $\eta_{s,t}(x)$  be a solution of the following stochastic differential equation:

$$\eta_{s,t}(x) = x + \int_s^t A(r, \eta_{s,r}(x)) d\beta(r) + \int_s^t B(r, \eta_{s,r}(x)) dr,$$

where  $\beta(t)$  is the standard  $E'$ -Wiener process. By the assumptions (H1) and (H2), if  $p \geq p_0$  and  $x \in E'_p$ , then the solution of the above equation is uniquely obtained by the usual method of successive approximations in  $E'_p$ .

We will assume the following condition:

(H4)  $(L(t)F)(x)$  and  $(U(t,s)F)(x) = E[F(\eta_{s,t}(x))] \in \mathfrak{H}_E$ , if  $F \in \mathfrak{H}_E$ .

Let  $W_F(t)$ ,  $F \in \mathfrak{H}_E$ , be the  $\mathcal{L}(\mathfrak{H}_E)$ -Wiener process and  $L(t)$  the diffusion

operator defined above. Then we will prove

Proposition 1. Under the assumptions (H1)-(H4) the continuous  $\mathcal{L}(\mathfrak{H}_E)$ -process solution of (1.1) such that for some  $0 < \alpha < 1$ ,  $E[|X_F(0)|^{2+\alpha}] < \infty$  is uniquely given as follows:

$$X_F(t) = X_{U(t,0)}F(0) + W_F(t) + \int_0^t W_L(s)U(t,s)F(s)ds.$$

Proof. Under the assumptions (H1)-(H4),  $L(t)$  is a continuous linear operator from  $\mathfrak{H}_E$  into itself. We use the following lemma which will be proved later.

Lemma 1. Suppose that the conditions (H1)-(H4) hold. Then  $L(t)$  generates the Kolmogorov evolution operator  $U(t,s)$  from  $\mathfrak{H}_E$  into itself such that

- (1)  $U(t,s)$  is a continuous linear operator from  $\mathfrak{H}_E$  into itself,
- (2) for any  $F \in \mathfrak{H}_E$ ,  $U(t,s)F$  is continuous from  $\{(t,s); 0 \leq s \leq t\}$  into  $\mathfrak{H}_E$ ,
- (3)  $U(t,t) = U(s,s) = \text{identity operator}$ ,
- (4)  $\frac{d}{dt} U(t,s)F = U(t,s)L(t)F$ ,  $0 \leq s \leq t$  on  $\mathfrak{H}_E$ ,
- (5)  $\frac{d}{ds} U(t,s)F = -L(s)U(t,s)F$ ,  $0 \leq s \leq t$ ,  $t > 0$  on  $\mathfrak{H}_E$ .

Further for any integers  $p \geq p_0$ ,  $q \geq 0$ ,  $n \geq 0$ ,  $j \geq 1$  and any  $T > 0$  and  $F \in \mathfrak{H}_E$ , we have

$$(2.1) \quad \|U(\cdot, s')F - U(t, s)F\|_{p,q,n}^{2j} \leq C_5(T, F, p, q, n) \{ |t-t'|^j + |s-s'|^j \},$$

$$0 \leq s, t, s', t' \leq T.$$

First we will verify that the integral in Proposition 1 is well defined by showing that for any fixed  $F \in \mathfrak{H}_E$ ,  $W_L(s)U(t,s)F(s)$  is continuous in  $(t,s)$ . Since  $W_F(t)$  is a Gaussian additive process with mean 0 and variance  $V_t(F)$ , we get for any integer  $n \geq 1$ ,

$$(2.2) \quad E[|W_F(t_1) - W_F(t_2)|^{2n}] \leq C_6(T)(V_{t_1}(F) - V_{t_2}(F))^n, \quad 0 \leq t_1, t_2 \leq T.$$

We choose an integer  $k \geq 4$  such that  $2k\lambda(m_1, q_1, T) > 2$ , where  $m_1$  and  $q_1$  are the numbers which appeared in (1.7) and  $\lambda(m_1, q_1, T)$  is the number in (H3). For  $0 \leq s, t, s', t' \leq T$ , the inequalities (2.1) and (2.2) yield, together with (H3),

$$(2.3) \quad E\{|W_{L(s)}U(t, s)F(s') - W_{L(s)}U(t, s)F(s)|^{2k}\} \\ \leq C_7(T)(V_s(L(s)U(t, s)F) - V_s(L(s)U(t, s)F))^k$$

and

$$(2.4) \quad E\{|W_{L(s')}U(t', s')F(s') - W_{L(s)}U(t, s)F(s')|^{2k}\} \\ \leq C_8(T)\|L(s')U(t', s')F - L(s)U(t, s)F\|_{p_1, q_1, m_1}^{2k} \\ \leq C_9(T)\{\|U(t', s')F - U(t, s)F\|_{p_1, q_1, m_1+1}^{2k} + \|U(t', s')F - U(t, s)F\|_{p_1, q_1, m_1+2}^{2k} \\ + |s' - s|^{2k\lambda(m_1, q_1, T)}\} \\ \leq C_{10}(T)\{|t - t'|^k + |s - s'|^k + |s' - s|^{2k\lambda(m_1, q_1, T)}\}.$$

The inequalities (2.3) and (2.4) are sufficient for the Kolmogorov-Totoki criterion [25] for continuity in  $(t, s)$ . The continuity of  $W_{L(s)}U(t, s)L(t)F(s)$  in  $(t, s)$  can be proved similarly.

Now we proceed to the proof of the existence of solutions for (1.1). Taking the relation  $U(t, s)F = F + \int_s^t U(\tau, s)L(\tau)F d\tau$ , the continuity of  $W_{L(s)}U(\tau, s)L(\tau)F(s)$  in  $\tau$ , the linearity of  $W_*(s)$  and the  $L^2$ -continuity of  $W_*(s)$ , into account, we have

$$W_{L(s)}U(t, s)F(s) = W_{L(s)}F(s) + W_{L(s)}\int_s^t U(\tau, s)L(\tau)F d\tau(s) \\ = W_{L(s)}F(s) + \int_s^t W_{L(s)}U(\tau, s)L(\tau)F(s) d\tau.$$

so that by making use of the continuity of  $W_{L(s)U(\tau,s)L(\tau)F(s)}$  in  $(\tau,s)$  again, we get

$$\begin{aligned}
 (2.5) \quad & \int_0^t W_{L(s)U(t,s)F(s)} ds \\
 &= \int_0^t W_{L(s)F(s)} ds + \int_0^t \left( \int_s^t W_{L(s)U(\tau,s)L(\tau)F(s)} d\tau \right) ds \\
 &= \int_0^t W_{L(s)F(s)} ds + \int_0^t \left( \int_0^\tau W_{L(s)U(\tau,s)L(\tau)F(s)} ds \right) d\tau \\
 &= \int_0^t (W_{L(\tau)F(\tau)} + \int_0^\tau W_{L(s)U(\tau,s)L(\tau)F(s)} ds) d\tau \\
 &= \int_0^t (X_{L(\tau)F(\tau)} - X_{U(\tau,0)L(\tau)F(0)}) d\tau.
 \end{aligned}$$

Combining the  $L^2$ -continuity of  $X_F(0)$  in the definition of  $\mathcal{Y}(\mathfrak{H}_E)$ -process and the Jensen inequality such that  $E[|X_F(0)|^{2+\alpha}] \leq E[|X_F(0)|^2]^\alpha$ , we get that  $E[|X_F(0)|^{2+\alpha}]$  is continuous in  $\mathfrak{H}_E$ . Hence there exist positive integers  $p_2 \geq p_0, q_2$  and  $m_2$  such that

$$(2.6) \quad E[|X_F(0)|^{2+\alpha}] \leq C_{11} \|F\|_{p_2, q_2, m_2}^{2+\alpha}.$$

Therefore the Kolmogorov criterion for continuity, together with the inequalities (2.1) in Lemma 1 and (2.6), yields the continuity of  $X_{U(\tau,0)L(\tau)F(0)}$  in  $\tau$ . Thus we get

$$(2.7) \quad \int_0^t X_{U(\tau,0)L(\tau)F(0)} d\tau = X_{U(t,0)F(0)} - X_F(0).$$

The equalities (2.5) and (2.7) show that  $X_F(t)$  is a solution of the equation (1.1).

Following H. Komatsu [12], we now prove the uniqueness of  $L^2$ -continuous solutions for the equation (1.1). Let  $Y_1(t,F)$  and  $Y_2(t,F)$  be the two continuous  $\mathcal{Y}(\mathfrak{H}_E)$ -process solutions for the equation (1.1). First we remark by the Baire category theorem that for each  $T > 0$ , we have some natural number



$p_3 \geq p_0, q_3$  and  $m_3$  such that

$$(2.8) \quad \max_{i=1,2} \sup_{0 \leq t \leq T} E[Y_i(t, F)^2] \leq C_{12}(T) \|F\|_{p_3, q_3, m_3}.$$

Define  $v(t, F) = Y_1(t, F) - Y_2(t, F)$ . Then for any  $a > 0$ , we will prove  $\frac{d}{dt} E[v(t, U(a, t)F)^2] = 0$  for  $t \in (0, a]$ . The inequality (2.8) and the strong continuity of  $U(t, s)$ , ((2) in Lemma 1), yield

$$\begin{aligned} & E\left[\left|\frac{v(s, U(a, s)F)^2 - v(t, U(a, t)F)^2}{s - t}\right|\right] \\ & \leq C_{13}(T, F) E\left[\left(\frac{v(s, U(a, s)F) - v(t, U(a, t)F)}{s - t}\right)^2\right]^{1/2}, \quad s, t \in (0, a] \subset [0, T]. \end{aligned}$$

The inequality (2.8) and the strong continuity of  $L(t)$  and  $U(t, s)$  imply that

$$(2.9) \quad \lim_{s \rightarrow t} E\left[\left|\frac{v(s, U(a, t)F) - v(t, U(a, t)F)}{s - t} - v(t, L(t)U(a, t)F)\right|^2\right] = 0.$$

By the strong continuity of  $U(t, s)$ , we get similarly

$$(2.10) \quad \begin{aligned} \lim_{s \rightarrow t} E\left[\left|\frac{v(s, [U(a, s) - U(a, t)]F) - v(t, [U(a, s) - U(a, t)]F)}{s - t} \right. \right. \\ \left. \left. - v(t, L(t)[U(a, s) - U(a, t)]F)\right|^2\right] = 0. \end{aligned}$$

Since  $L(t)$  generates the Kolmogorov evolution operator  $U(t, s)$ , we have

$$\begin{aligned} \lim_{s \rightarrow t} E\left[\left|v(t, L(t)U(a, s)F) - v(t, L(t)U(a, t)F)\right|^2\right] &= 0 \\ \lim_{s \rightarrow t} E\left[\left|v(t, L(t)U(a, t)F) + v(t, \frac{U(a, s) - U(a, t)}{s - t}F)\right|^2\right] &= 0. \end{aligned}$$

so that we get

$$(2.11) \quad \lim_{s \rightarrow t} E\left[\left|v(t, L(t)U(a, s)F) + \frac{v(t, U(a, s)F) - v(t, U(a, t)F)}{s - t}\right|^2\right] = 0.$$

From (2.9), (2.10) and (2.11), we get the desired equality claimed above.

Hence  $E[v(t, U(a, t)F)^2] = \text{constant}$ . Then letting  $t \rightarrow 0$ , by (2.8) and the definition of continuity of an  $\mathcal{L}(\mathfrak{D}_E)$ -process in  $t$ , we have the constant = 0. Taking the equalities  $E[v(t, U(a, t)F)^2] = E[(v(t, F) + v(t, [U(a, t) - U(a, a)]F))^2]$  and  $\lim_{t \rightarrow a} E[v(t, [U(a, t) - U(a, a)]F)^2] = 0$ , into account, we have  $E[v(a, F)^2] = 0$  for any  $a > 0$ , which implies  $v(a, F) = 0$  almost surely. Thus the proof is completed.

### §3. Proof of Lemma 1.

As in [20], [21], we will treat the generation problem via the stochastic method.

For any  $F$  in  $\mathfrak{D}_E$ , we recall the definition of  $U(t, s)$ :

$$(U(t, s)F)(x) = E[F(\eta_{s, t}(x))].$$

To examine that  $U(t, s)$  is the evolution operator stated in Lemma 1, we will check some regularities and integrabilities for  $\eta_{s, t}(x)$ . It is obvious that if  $p \geq p_0$  and  $x \in E'_p$ ,  $\eta_{s, t}(x) \in E'_p$ , so that for  $h \in E'_{p_4}$ ,  $\eta_{s, t}(x+h) \in E'_{p_5}$ , where  $p_5 = p \vee p_4$ . Here  $a \vee b = \max\{a, b\}$ . Following Kunita (p. 219 of [13]), we will show that  $\xi_{s, t}(\tau) := \frac{1}{\tau}(\eta_{s, t}(x+\tau h) - \eta_{s, t}(x))$  has a continuous extension at  $\tau = 0$  for any  $s, t$  a.s. in  $E'_{p_5}$ . This can be proved by appealing to the Kolmogorov-Totoki criterion for continuity [25].

Lemma 2. For any  $T > 0$  and any integer  $j \geq 1$ , we have

$$E[\|\xi_{s, t}(\tau) - \xi_{s', t'}(\tau')\|_{-p_5}^{2j}] \leq C_{14}(T, h)\{|s-s'|^j + |t-t'|^j + |\tau-\tau'|^j\},$$

$$0 \leq s, s', t, t', \tau, \tau' \leq T.$$

First we will show the following inequality. Let  $A(r)$  be a well measurable random linear operator from  $E'_1$  to  $E'_{p_0}$  such that

$E[\int_s^t \|A(r)\|_2^2 dr] < \infty$ . Then we have

Lemma 3. For any integer  $j \geq 1$ ,

$$E[\|\int_s^t A(r) d\beta(r)\|_{-p_0}^{2j}] \leq C_{15}(j) E[(\int_s^t \|A(r)\|_2^2 dr)^j].$$

Proof. Let  $(\cdot, \cdot)_{-p_0}$  be the inner product in  $E'_{p_0}$  such that  $(x, x)_{-p_0} = \|x\|_{-p_0}^2$ . Setting  $\theta(x) = (x, x)_{-p_0}^j$  and  $y(t) = \int_s^t A(r) d\beta(r)$  and applying the Itô formula, (Kuo [15]), for  $\theta(y(t))$ , we get

$$\begin{aligned} (3.1) \quad E[\|y(t)\|_{-p_0}^{2j}] &= \frac{1}{2} E[\int_s^t \text{trace}_{E_0} D^2 \theta(y(r)) \circ [A(r) \times A(r)] dr] \\ &= \frac{1}{2} E[\int_s^t \sum_{i=1}^{\infty} \{2^{2j(j-1)} (y(r), A(r) h_i^{(0)})_{-p_0}^2 \|y(r)\|_{-p_0}^{2(j-2)} \\ &\quad + 2j \|A(r) h_i^{(0)}\|_{-p_0}^2 \|y(r)\|_{-p_0}^{2(j-1)}\} dr] \\ &\leq (j+2j(j-1)) E[\int_s^t \|A(r)\|_2^2 \|y(r)\|_{-p_0}^{2(j-1)} dr]. \end{aligned}$$

By Hölder's inequality and the martingale inequality, the right hand side of (3.1) is dominated by

$$\begin{aligned} &(j+2j(j-1)) E[\sup_{s \leq r \leq t} \|y(r)\|_{-p_0}^{2j}]^{j-1/j} E[(\int_s^t \|A(r)\|_2^2 dr)^j]^{1/j} \\ &\leq (2j^2-j)(2j/(2j-1))^{2(j-1)} E[\|y(t)\|_{-p_0}^{2j}]^{j-1/j} E[(\int_s^t \|A(r)\|_2^2 dr)^j]^{1/j}, \end{aligned}$$

which completes the proof of Lemma 3.

Proof of Lemma 2. Now for the convenience of notations we will write  $dt = d\beta_0(t)$ ,  $d\beta(t) = d\beta_1(t)$ ,  $A_0(t, x) = B(t, x)$ ,  $A_1(t, x) = A(t, x)$ ,  $\|\cdot\|_0 = \|\cdot\|_{-p_0}$  and  $\|\cdot\|_1 = \|\cdot\|_2$ . Without loss of generality, we may assume

$0 \leq s < s' < t < t' \leq T$ . Then  $\xi_{s,t}(\tau) - \xi_{s',t}(\tau')$  is a sum of the following terms:

$$(3.2) \quad \sum_k \int_s^{s'} (\int_0^1 DA_k(r, \xi_{s,r}(\tau, y)) (\xi_{s,r}(\tau)) dy) d\beta_k(r),$$

where  $\xi_{s,r}(\tau, y) = \eta_{s,r}(x) + y(\eta_{s,r}(x+\tau h) - \eta_{s,r}(x))$ .

$$(3.3) \quad \sum_k \int_s^t (\int_0^1 DA_k(r, \xi_{s,r}(\tau, y)) (\xi_{s,r}(\tau)) - DA_k(r, \xi_{s',r}(\tau', y)) (\xi_{s',r}(\tau'))) dy) d\beta_k(r).$$

By Lemma 3 and the assumption (H2), the expectation of the  $2j$ -th power of the  $\|\cdot\|_{-p_5}$ -norm of (3.2) is dominated by

$$C_{16} \sum_k E[(\int_s^{s'} (\int_0^1 DA_k(r, \xi_{s,r}(\tau, y)) (\xi_{s,r}(\tau)) dy)^2 dr)^j] \\ \leq C_{17} \sum_k |s'-s|^{j-1} E[\int_s^{s'} \|\xi_{s,r}(\tau)\|_{-p_5}^{2j} dr].$$

Again using Lemma 3, assumption (H2) and the Gronwall lemma, we have

$$(3.4) \quad E[\|\eta_{s,t}(x) - \eta_{s,t}(y)\|_{-p_5}^{2j}] \leq C_{18} \|x-y\|_{-p_5}^{2j}, \quad x, y \in E'_{p_5},$$

which implies

$$(3.5) \quad E[\int_s^{s'} \|\xi_{s,r}(\tau)\|_{-p_5}^{2j} dr] \leq C_{18} \|h\|_{-p_5}^{2j} |s'-s|.$$

Since the integrand in (3.3)

$$= \int_0^1 DA_k(r, \xi_{s,r}(\tau, y)) (\xi_{s,r}(\tau) - \xi_{s',r}(\tau')) dy \\ + \int_0^1 (\int_0^1 D^2 A_k(r, \gamma_{s,s',r}(\tau, \tau', y_1)) (\xi_{s,r}(\tau, y) - \xi_{s',r}(\tau', y)) dy_1) (\xi_{s',r}(\tau')) dy,$$

where  $\gamma_{s,s',r}(\tau, \tau', y_1) = \xi_{s',r}(\tau', y) + y_1(\xi_{s,r}(\tau, y) - \xi_{s',r}(\tau', y))$ , the

$\|\cdot\|_k$ -norm of the integrand is dominated by

$$(3.6) \quad C_{19} \{ \|\xi_{s,r}(\tau) - \xi_{s',r}(\tau')\|_{-p_5} + (\|\eta_{s,r}(x) - \eta_{s',r}(x)\|_{-p_5} + \|\eta_{s,r}(x+\tau h) - \eta_{s',r}(x+\tau'h)\|_{-p_5}) \|\xi_{s',r}(\tau')\|_{-p_5} \}.$$

By Lemma 3 and (3.6), the expectation of the  $2j$ -th power of  $\|\cdot\|_{-p_5}$ -norm of (3.3) is dominated by

$$(3.7) \quad C_{20} \{ \int_s^t E[\|\xi_{s,r}(\tau) - \xi_{s',r}(\tau')\|_{-p_5}^{2j}] dr + \int_s^t E[\|\eta_{s,r}(x) - \eta_{s',r}(x)\|_{-p_5}^{4j}]^{1/2} E[\|\xi_{s',r}(\tau')\|_{-p_5}^{4j}]^{1/2} dr + \int_s^t E[\|\eta_{s,r}(x+\tau h) - \eta_{s',r}(x+\tau'h)\|_{-p_5}^{4j}]^{1/2} E[\|\xi_{s',r}(\tau')\|_{-p_5}^{4j}]^{1/2} dr \}.$$

From the assumptions (H1) and (H2), we get

$$\|A_k(r, \eta_{s,r}(x)) - A_k(r, \eta_{s',r}(x'))\|_k \leq C_{21} \|\eta_{s,r}(x) - \eta_{s',r}(x')\|_{-p_5}$$

and taking the expectations of the  $2n$ -th power of both sides of  $\|\cdot\|_{-p_5}$ -norm of the following inequality:

$$\begin{aligned} & \|\eta_{s,t}(x) - \eta_{s',t'}(x')\|_{-p_5} \\ & \leq \|\sum_k \int_s^{s'} A_k(r, \eta_{s,r}(x)) d\beta_k(r)\|_{-p_5} \\ & + \|\sum_k \int_t^{t'} A_k(r, \eta_{s',r}(x')) d\beta_k(r)\|_{-p_5} \\ & + \|\sum_k \int_s^t (A_k(r, \eta_{s,r}(x)) - A_k(r, \eta_{s',r}(x')) d\beta_k(r)\|_{-p_5}. \end{aligned}$$

we have, by Lemma 3, similarly

$$E[\|\eta_{s,t}(x) - \eta_{s',t'}(x')\|_{-p_5}^{2n}] \\ \leq C_{22}(T)\{|t-t'|^n + |s-s'|^n + \int_s^t E[\|\eta_{s,r}(x) - \eta_{s',r}(x')\|_{-p_5}^{2n}]dr\}.$$

Noticing that  $\eta_{s,r}(x) = \eta_{s',r}(\eta_{s,s}(x))$  and  $\eta_{s',r}(\cdot)$  is independent of  $\eta_{s,s}(\cdot)$  and using (3.4), we get

$$E[\|\eta_{s,r}(x) - \eta_{s',r}(x')\|_{-p_5}^{2n}] = \int_{E'} E[\|\eta_{s',r}(y) - \eta_{s',r}(x')\|_{-p_5}^{2n}] P(\eta_{s,s}(x) \in dy) \\ \leq \int_{E'} C_{23} \|y - x'\|_{-p_5}^{2n} P(\eta_{s,s}(x) \in dy) \\ = C_{23} E[\|\eta_{s,s}(x) - x'\|_{-p_5}^{2n}] \\ \leq C_{24} \{\|x - x'\|_{-p_5}^{2n} + |s' - s|^n\},$$

where  $P(\cdot)$  denotes the fundamental probability measure associated with  $\beta(t)$ .

Hence we obtain

$$(3.8) \quad E[\|\eta_{s,t}(x) - \eta_{s',t'}(x')\|_{-p_5}^{2n}] \leq C_{25}(T)\{|t-t'|^n + |s-s'|^n + \|x-x'\|_{-p_5}^{2n}\}.$$

Combining (3.2), (3.3), (3.4), (3.5), (3.7) and (3.8), we have

$$E[\|\xi_{s,t}(\tau) - \xi_{s',t'}(\tau')\|_{-p_5}^{2j}] \\ \leq C_{26}(T) \|h\|_{-p_5}^{2j} \{|t-t'|^j + |s-s'|^j + |\tau-\tau'|^{2j} \|h\|_{-p_5}^{2j}\}.$$

This completes the proof of Lemma 2.

Let  $\tau$  tend to 0, we have for each  $x \in E'_p$ ,

$$(3.9) \quad D\eta_{s,t}(x)(h) = h + \sum_k \int_s^t DA_k(r, \eta_{s,r}(x))(D\eta_{s,r}(x)(h)) d\beta_k(r).$$

For the higher order differentiations, the formula similar to (3.9) can be proved inductively, together with the following lemma.

Lemma 4. Suppose that a natural number  $q \geq p_0$  and any  $T > 0$ . Then for  $0 \leq s, t, s', t' \leq T$ , a natural number  $j$  and  $x, x', h_i \in E'_q$ ,  $i=1, 2, \dots, n$ , we have

$$(3.10) \quad E[\|D^n \eta_{s,t}(x)(h_1, h_2, \dots, h_n)\|_{-q}^{2j}] \leq C_{27}(T) \|h_1\|_{-q}^{2j} \|h_2\|_{-q}^{2j} \dots \|h_n\|_{-q}^{2j}.$$

$$(3.11) \quad E[\|D^n \eta_{s,t}(x)(h_1, h_2, \dots, h_n) - D^n \eta_{s',t'}(x')(h_1, h_2, \dots, h_n)\|_{-q}^{2j}] \\ \leq C_{28}(T) \{ |t-t'|^j + |s-s'|^j + \|x-x'\|_{-q}^{2j} \} \|h_1\|_{-q}^{2j} \|h_2\|_{-q}^{2j} \dots \|h_n\|_{-q}^{2j}.$$

Proof. First we will show (3.10) for the case  $n=1$ . By the assumptions (H1) and (H2), we get

$$\|DA_k(r, \eta_{s,r}(x))(D\eta_{s,r}(x)(h))\|_k \leq C_{29} \|D\eta_{s,r}(x)(h)\|_{-q},$$

so that taking the expectations of  $2j$ -th powers of  $\|\cdot\|_{-q}$  norms of both sides of (3.9) and using Lemma 3, we get

$$E[\|D\eta_{s,t}(x)(h)\|_{-q}^{2j}] \leq C_{30}(T) \{ \|h\|_{-q}^{2j} + \int_s^t E[\|D\eta_{s,r}(x)(h)\|_{-q}^{2j}] dr \}$$

and the Gronwall inequality gives (3.10) for the case where  $n=1$ . For  $n \geq 2$ , we will prove the inequality by mathematical induction. For  $h_1, h_2, \dots, h_n \in E'_q$ ,

$$(D^n \eta_{s,t}(x))(h_1, h_2, \dots, h_n) = \sum_k \int_s^t D^n(A_k(r, \eta_{s,r}(x)))(h_1, h_2, \dots, h_n) d\beta_k(r).$$

Since

$$(3.12) \quad D^n(A_k(r, \eta_{s,r}(x)))(h_1, h_2, \dots, h_n) \\ = DA_k(r, \eta_{s,r}(x))(D^n \eta_{s,r}(x)(h_1, h_2, \dots, h_n)) \\ + \text{finite sum of terms of the type}$$

$$(D^m A_k(r, \eta_{s,r}(x))) (D^{n_1} \eta_{s,r}(x) (h_{j_1(1)}, h_{j_2(1)}, \dots, h_{j_{n_1}(1)}), \\ D^{n_2} \eta_{s,r}(x) (h_{j_1(2)}, h_{j_2(2)}, \dots, h_{j_{n_2}(2)}), \dots, D^{n_m} \eta_{s,r}(x) (h_{j_1(m)}, h_{j_2(m)}, \dots, h_{j_{n_m}(m)})).$$

where  $2 \leq m \leq n$ ,  $n_1 + n_2 + \dots + n_m = n$  and  $0 \leq n_1 \leq n-1$ , so that using the inductive assumption, we get (3.10) by the same argument as before.

Before proceeding to the proof of (3.11), we note that for  $h \in E'_q$ ,  $\|D\eta_{s,t}(x)(h) - D\eta_{s',t'}(x')(h)\|_{-q}$  is dominated by

$$(3.13) \quad \sum_k \|\int_s^{s'} D(A_k(r, \eta_{s,r}(x)))(h) d\beta_k(r)\|_{-q} \\ + \sum_k \|\int_t^{t'} D(A_k(r, \eta_{s',r}(x')))(h) d\beta_k(r)\|_{-q} \\ + \sum_k \|\int_s^{s'} \{D(A_k(r, \eta_{s,r}(x)))(h) - D(A_k(r, \eta_{s',r}(x')))(h)\} d\beta_k(r)\|_{-q}.$$

Now, by the assumptions (H1) and (H2), we have

$$(3.14) \quad \|D(A_k(r, \eta_{s,r}(x)))(h) - D(A_k(r, \eta_{s',r}(x')))(h)\|_k \\ \leq \| \{DA_k(r, \eta_{s,r}(x)) - DA_k(r, \eta_{s',r}(x'))\} (D\eta_{s,r}(x)(h)) \|_k \\ + \|DA_k(r, \eta_{s',r}(x'))(D\eta_{s,r}(x)(h) - D\eta_{s',r}(x')(h))\|_k \\ \leq C_{31}(T) \{\|\eta_{s,r}(x) - \eta_{s',r}(x')\|_{-q} \|D\eta_{s,r}(x)(h)\|_{-q} \\ + \|D\eta_{s,r}(x)(h) - D\eta_{s',r}(x')(h)\|_{-q}\}.$$

Hence from (3.8), (3.13) and (3.14) we have

$$E[\|D\eta_{s,t}(x)(h) - D\eta_{s',t'}(x')(h)\|_{-q}^{2j}]$$



$$\leq C_{32}(T) \{ (|t-t'|^J + |s-s'|^J + \|x-x'\|_{-q}^{2J}) \|h\|_{-q}^{2J} \\ + \int_s^t E[\|D\eta_{s,t}(x)(h) - D\eta_{s',t'}(x')(h)\|_{-q}^{2J}] dr \},$$

which gives (3.11) by the Gronwall lemma for the case  $n=1$ . By (3.12) and the estimation of  $\|D^n \eta_{s,t}(x)(h_1, h_2, \dots, h_n) - D^n \eta_{s',t'}(x')(h_1, h_2, \dots, h_n)\|_{-q}$  similar to that in (3.13), mathematical induction and the Gronwall lemma yield the proof of (3.11) for  $n \geq 2$ .

For the proof of the generation problem of  $L(t)$  we proceed as follows. By the assumptions (H1) and (H2), (3.8) and (3.10) of Lemma 4, we may exchange the order of the differentiation and the integration. Then by the Itô formula [15], we have the pointwise Kolmogorov forward and backward equations as in the finite dimensional case (Theorem 1 (page 73) of [7]):

$$\frac{d}{dt} (U(t,s)F)(x) = (U(t,s)L(t)F)(x)$$

$$\frac{d}{ds} (U(t,s)F)(x) = -(L(s)U(t,s)F)(x).$$

Let  $p \geq 0$ ,  $q \geq 0$  and  $n \geq 0$  be integers and  $x \in E'_p$ . Since  $D^n(F(\eta_{s,t}(x)))(h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_n}^{(q)})$  is a finite sum of terms of the type

$$I = D^m F(\eta_{s,t}(x)) (D^{n_1} \eta_{s,t}(x)(h_{j_1(1)}^{(q)}, h_{j_2(1)}^{(q)}, \dots, h_{j_{n_1}(1)}^{(q)}), D^{n_2} \eta_{s,t}(x) \\ (h_{j_1(2)}^{(q)}, h_{j_2(2)}^{(q)}, \dots, h_{j_{n_2}(2)}^{(q)}), \dots, D^{n_m} \eta_{s,t}(x)(h_{j_1(m)}^{(q)}, h_{j_2(m)}^{(q)}, \dots, h_{j_{n_m}(m)}^{(q)})),$$

$$n_1 + n_2 + \dots + n_m = n,$$

so that from the nuclearity of  $E$  and (3.10), we have an integer

$q' > \max\{p, p_0, q\}$  such that

$$(3.15) \quad \sum_{j=1}^{\infty} \|h_j^{(q)}\|_{-q}^2 < +\infty$$

and

$$(3.16) \quad E[|I|^2] \leq \|F\|_{q', q', n}^2 E[e^{2\|\eta_{s,t}(x)\|_{-q'}} \|D^{n_1} \eta_{s,t}(x)(h_{j_1(1)}^{(q)}, h_{j_2(1)}^{(q)}, \dots, h_{j_{n_1}(1)}^{(q)})\|_{-q'}^2 \cdot \dots \cdot \|D^{n_2} \eta_{s,t}(x)(h_{j_1(2)}^{(q)}, h_{j_2(2)}^{(q)}, \dots, h_{j_{n_2}(2)}^{(q)})\|_{-q'}^2 \cdot \dots \cdot \|D^{n_m} \eta_{s,t}(x)(h_{j_1(m)}^{(q)}, h_{j_2(m)}^{(q)}, \dots, h_{j_{n_m}(m)}^{(q)})\|_{-q'}^2]$$

$$\leq C_{33} \|F\|_{q', q', n}^2 \|h_{i_1}^{(q)}\|_{-q}^2 \|h_{i_2}^{(q)}\|_{-q}^2 \dots \|h_{i_n}^{(q)}\|_{-q}^2 E[e^{4\|\eta_{s,t}(x)\|_{-q'}}]^{1/2}.$$

Here we will prove

Lemma 5. For any  $\alpha > 0$  and  $T > 0$ , there exists a constant  $C_{34} = C_{34}(\alpha, T)$  such that

$$\sup_{0 \leq s, t \leq T} E[e^{\alpha \|\eta_{s,t}(x)\|_{-q'}}] \leq C_{34} e^{\alpha \|x\|_{-q'}}.$$

Proof. By (H1),  $\|\eta_{s,t}(x)\|_{-q} \leq \|x\|_{-q} + C_{35} + \|\int_s^t \Lambda(r, \eta_{s,r}(x)) d\beta(r)\|_{-q}$ .

Following [8], it is enough to prove  $E[\exp(\|\int_s^t \alpha \Lambda(r, \eta_{s,r}(x)) d\beta(r)\|_{-q})] \leq C_{36}$ .

Setting  $y_{s,t}(x) = \int_s^t \alpha \Lambda(r, \eta_{s,r}(x)) d\beta(r)$ , by the Itô formula and the assumption (H1), we get for any integer  $m \geq 2$ ,

$$(3.17) \quad E[\|y_{s,t}(x)\|_{-q}^m] \leq E[(1 + \|y_{s,t}(x)\|_{-q}^2)^{m/2}]$$

$$\leq E[1 + \int_s^t \frac{1}{2} (2^{\frac{m}{2}} (1 + \|y_{s,r}(x)\|_{-q}^2)^{\frac{m}{2}-1} \alpha^2 \|\Lambda(r, \eta_{s,r}(x))\|_2^2]$$

$$\begin{aligned}
& + 4^{\frac{m}{2}} \left( \frac{m}{2} - 1 \right) (1 + \|y_{s,r}(x)\|_{-q}^2)^{\frac{m}{2} - 2} \alpha^2 \left( \sum_{i=1}^{\infty} (y_{s,r}(x), A(r, \eta_{s,r}(x)) h_i^{(0)})_{-q}^2 \right) dr \\
& \leq 1 + 2 \left( \frac{m}{2} \right)^2 \alpha^2 C_{37} \int_s^t E \left[ (1 + \|y_{s,r}(s)\|_{-q}^2)^{\frac{m}{2} - 1} \right] dr,
\end{aligned}$$

where  $C_{37} = \sup_{\substack{x \in E \\ 0 \leq t \leq T}} \|A(t, x)\|_2^2$  and  $\|x\|_{-q}^2 = (x, x)_{-q}$ . If we use (3.17)

recursively, the rest is similar to the argument in [8], which completes the proof.

Therefore (3.15), (3.16) and Lemma 5 yield

$$\|U(t, s)F\|_{p, q, n} \leq C_{38}(T) \|F\|_{q', q', n}, \quad t, s \in [0, T],$$

which implies that  $U(t, s)$  is a continuous linear operator from  $\mathfrak{D}_E$  into itself.

In the same way as in [21], if we prove the strong continuity of  $U(t, s)F$  in  $(t, s)$ , the pointwise Kolmogorov forward and backward equations imply that  $L(t)$  generates the evolution operator  $U(t, s)$ . Since  $\|U(t, s) - U(t', s')\|_{p, q, n}^{2j}$  is dominated by a finite sum of terms of the type

$$\begin{aligned}
& \sup_{\substack{x \in E \\ p}} e^{-2j\|x\|_p} \sum_{\substack{j_1^{(1)}, j_2^{(1)}, \dots, j_{m_1}^{(1)} \\ \vdots \\ j_1^{(m)}, j_2^{(m)}, \dots, j_{n_m}^{(m)}}} E[|D^m F(\eta_{s,t}(x))| (D^{n_1} \eta_{s,t}(x) \\
& (h_{j_1^{(1)}}^{(q)}, h_{j_2^{(1)}}^{(q)}, \dots, h_{j_{n_1}^{(1)}}^{(q)}), D^{n_2} \eta_{s,t}(x) (h_{j_1^{(2)}}^{(q)}, h_{j_2^{(2)}}^{(q)}, \dots, h_{j_{n_2}^{(2)}}^{(q)}), \dots, \\
& D^{n_m} \eta_{s,t}(x) (h_{j_1^{(m)}}^{(q)}, h_{j_2^{(m)}}^{(q)}, \dots, h_{j_{n_m}^{(m)}}^{(q)}) - D^m F(\eta_{s',t'}(x)) (D^{n_1} \eta_{s',t'}(x)
\end{aligned}$$

$$\begin{aligned}
& (h_{j_1(1)}^{(q)}, h_{j_2(1)}^{(q)}, \dots, h_{j_{n_1}(1)}^{(q)}) \cdot D_{\eta_{s', t'}^{n_2}}(x) (h_{j_1(2)}^{(q)}, h_{j_2(2)}^{(q)}, \dots, h_{j_{n_2}(2)}^{(q)}) \cdot \\
& \dots, D_{\eta_{s', t'}^{n_m}}(x) (h_{j_1(m)}^{(q)}, h_{j_2(m)}^{(q)}, \dots, h_{j_{n_m}(m)}^{(q)}) |^{2j}].
\end{aligned}$$

so that by (3.8), Lemmas 4 and 5 and the nuclearity of  $E$ , we have

$$\|U(t, s)F - U(t', s')F\|_{p, q, n}^{2j} \leq C_{39} \|F\|_{q', q', n+1}^{2j} \{|t-t'|^j + |s-s'|^j\}.$$

This completes the proof of Lemma 1.

#### §4. Generation of the Kolmogorov Evolution Operator

In this Section, we will discuss assumption (H4). Let  $K$  be a separable Hilbert space. We call a  $K$ -valued functional

$G(x) = g(\langle x, f_1 \rangle, \langle x, f_2 \rangle, \dots, \langle x, f_n \rangle)$ ,  $f_1, f_2, \dots, f_n \in E$ , a smooth functional if

$g(x): \mathbb{R}^n \rightarrow K$  is a  $C^\infty$ -function. Further we call  $G(x)$  a bounded smooth

functional if  $g(x)$  itself and all the derivatives of  $g(x)$  are bounded. The

coefficients  $A(t, x)$  and  $B(t, x)$  are said to be approximated by bounded smooth

functionals on  $E'$  if for any integers,  $p \geq p_0$ ,  $q \geq 0$  and  $n \geq 0$ , there exist

sequences of bounded smooth functionals

$$A_m(t, x) = a_m(t, \langle x, f_1 \rangle, \langle x, f_2 \rangle, \dots, \langle x, f_{k_m} \rangle)$$

and

$$B_m(t, x) = b_m(t, \langle x, f_1 \rangle, \langle x, f_2 \rangle, \dots, \langle x, f_{k_m} \rangle)$$

such that the following conditions are satisfied:

(4.1)  $A_m(t, x)$  and  $B_m(t, x)$  satisfy the conditions  $(H_1)$ ,  $(H_2)$  and  $(H_3)$ .

(4.2) For any  $T > 0$ ,

$$\lim_{m \rightarrow \infty} \sup_{\substack{x \in E'_p \\ 0 \leq t \leq T}} \|A(t, x) - A_m(t, x)\|_2^2 = 0,$$

$$\lim_{m \rightarrow \infty} \sup_{\substack{x \in E'_p \\ 0 \leq t \leq T}} \|B(t, x) - B_m(t, x)\|_{-p_0} = 0,$$

$$\lim_{m \rightarrow \infty} \sup_{\substack{x \in E'_p \\ 0 \leq t \leq T}} \|D^k A(t, x) - D^k A_m(t, x)\|_{H.S.}^{(q)} = 0, \quad k=1, 2, \dots, n,$$

$$\lim_{m \rightarrow \infty} \sup_{\substack{x \in E'_p \\ 0 \leq t \leq T}} \|D^k B(t, x) - D^k B_m(t, x)\|_{H.S.}^{(q)} = 0, \quad k=1, 2, \dots, n.$$

Proposition 2. Suppose that the coefficients  $A(t, x)$  and  $B(t, x)$  are approximated by bounded smooth functionals on  $E'$ . Then if  $F \in \mathcal{D}_E$ ,  
 $U(t, s)F(x) = E[F(\eta_{s, t}(x))] \in \mathcal{D}_E$ .

Proof. It is convenient to use the notation  $A_0(t, x) = B(t, x)$  and  $A_1(t, x) = A(t, x)$ . For any integers  $p \geq 0$ ,  $q \geq 0$  and  $n \geq 0$ , we choose an integer  $q' > \max\{p, p_0, q\}$  such that

$$(4.3) \quad \sum_{j=1}^{\infty} \|h_j^{(q)}\|_{-q}^2 < +\infty,$$

since  $E$  is a nuclear Fréchet space. Then by the assumptions, for any  $0 < \delta < 1$  and  $A_k(t, x)$ ,  $k=0, 1$ , there exist bounded smooth functionals  $\tilde{A}_k(t, x) = \tilde{a}_k(t, \langle x, \zeta_1 \rangle, \langle x, \zeta_2 \rangle, \dots, \langle x, \zeta_{m_k} \rangle)$ ,  $k=0, 1$  such that

$$(4.4) \quad \sum_{\ell=0}^{n+1} \sup_{\substack{x \in E'_{q'} \\ 0 \leq t \leq T}} \|D^\ell A_k(t, x) - D^\ell \tilde{A}_k(t, x)\|_{H.S.}^{(q')} < \delta.$$

For sufficiently large  $N$ , we put

$$z_{s,t}^N(x) = x + \sum_k \int_s^t \tilde{A}_k(t_1, x + \sum_k \int_s^{t_1} \tilde{A}_k(t_2, \dots \\ \dots, x + \sum_k \int_s^{t_{N-1}} \tilde{A}_k(t_N, x) d\beta_k(t_N)) \dots) d\beta_k(t_1).$$

Setting

$$\hat{z}_{s,t}^{(n)}(x) = x + \sum_k \int_s^t \tilde{A}_k(t_1, x + \sum_k \int_s^{t_1} \tilde{A}_k(t_2, \dots, x + \sum_k \int_s^{t_{n-1}} \tilde{A}_k(t_n, \eta_{s,t_n}(x)) d\beta_k(t_n)) \dots) d\beta_k(t_1), \\ n=1, 2, \dots, N, \text{ where } t_0=t, \text{ by Lemma 3, we have for any } x \in E'_p, 0 \leq s, t \leq T \text{ and} \\ \text{any integer } j \geq 1.$$

$$(4.5) \quad E[\|\eta_{s,t}(x) - z_{s,t}^N(x)\|_{-q}^{2j}] \\ \leq 2^{2j-1} E[\|\eta_{s,t}(x) - \hat{z}_{s,t}^{(1)}(x)\|_{-q}^{2j}] \\ + \sum_{k=2}^N (2^{2j-1})^k E[\|\hat{z}_{s,t}^{(k-1)}(x) - \hat{z}_{s,t}^{(k)}(x)\|_{-q}^{2j}] \\ + (2^{2j-1})^N E[\|\hat{z}_{s,t}^{(N)}(x) - z_{s,t}^N(x)\|_{-q}^{2j}] \\ \leq (2^{2j-1})^2 \delta^{2j} R_T + \sum_{k=2}^N (2^{2j-1})^{2k} 2^{2j(k-1)} \delta^{2j} R_T^k / k! \\ + (2^{2j-1})^{2N} R^N 2^{2j} M^{2jN} T^N / N! \\ \leq \delta^{2j} \exp(2^{2(2j-1)} R(MV_1)^{2j} T) + (2^{2j-1})^{2N} R^N 2^{2j} M^{2jN} T^N / N!,$$

where  $M = \max_k \max_{0 \leq \ell \leq n+1} \sup_{\substack{x \in E'_p \\ 0 \leq t \leq T}} \|\tilde{A}_k(t, x)\|_{H.S.}^{(q')}$  and  $R = C_{15}(j) T^{j-1} + T^{2j-1}$ . Hence

for any  $\epsilon > 0$ , if we take sufficiently small  $\delta$  and large  $N$ , we have

$$(4.6) \quad \sup_{x \in E'_p} E[\|\eta_{s,t}(x) - z_{s,t}^N(x)\|_{-q}^{2j}] < \epsilon.$$

Next we will verify by mathematical induction that for any integer

$1 \leq k \leq n$  and any  $\epsilon > 0$ , there exists an integer  $N(k, \epsilon)$  such that if  $N \geq N(k, \epsilon)$ ,

$$(4.7) \quad E[\|D_{s,t}^k \eta_{s,t}(x)(h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)}) - D_{s,t}^k z_{s,t}^N(x)(h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)})\|_{-q}^{2j}] < \epsilon.$$

For any  $\epsilon' > 0$ , (4.5) gives that if we take sufficiently small  $\delta$  and large  $n(\epsilon')$ , then for any  $N \geq n(\epsilon')$

$$(4.8) \quad \sup_{x \in E'_p} E[\|\eta_{s,t}(x) - z_{s,t}^N(x)\|_{-q}^{4j}]^{1/2} < \epsilon'.$$

Here we need the following lemma for later use. In a manner similar to that in the proofs of (3.10) and Lemma 5, we get

Lemma 6. For any integers  $q \geq p_0$ ,  $j \geq 1$ ,  $n \geq 1$  and any  $T > 0$ , we have

$$(4.9) \quad \sup_{0 \leq s, t \leq T} E[\|D_{s,t}^n z_{s,t}^N(x)(h_1, h_2, \dots, h_n)\|_{-q}^{2j}] \leq C_{40}(T) \|h_1\|_{-q}^{2j} \|h_2\|_{-q}^{2j} \dots \|h_n\|_{-q}^{2j}, \quad x, h_i, i = 1, 2, \dots, n \in E'_q.$$

For any  $\alpha > 0$  and  $T > 0$ ,

$$(4.10) \quad \sup_{0 \leq s, t \leq T} E[e^{\alpha \|z_{s,t}^N(x)\|_{-q}}] \leq C_{41} e^{\alpha \|x\|_{-q}}.$$

For any  $f \in E$  and any  $\alpha > 0$  and  $T > 0$ , there exists  $C_{42} = C_{42}(f, \alpha, T)$  such that

$$(4.11) \quad \sup_{0 \leq s, t \leq T} \max\{E[\exp(\alpha \sqrt{|\langle \eta_{s,t}(x), f \rangle|})], E[\exp(\alpha \sqrt{|\langle z_{s,t}^N(x), f \rangle|})]\} \leq C_{42} \exp(\alpha \sqrt{|\langle x, f \rangle|}).$$

Setting

$$\begin{aligned}
& y_{s,t}^{m,N}(x)(h_{i_1}^{(q)}) \\
&= h_{i_1}^{(q)} + \sum_k \int_s^t \tilde{D}\tilde{A}_k(t_1, z_{s,t_1}^{N-1}(x))(h_{i_1}^{(q)}) + \sum_k \int_s^{t_1} \tilde{D}\tilde{A}_k(t_2, z_{s,t_2}^{N-2}(x))(h_{i_1}^{(q)}) \\
&\quad + \dots + \sum_k \int_s^{t_{m-1}} \tilde{D}\tilde{A}_k(t_m, \eta_{s,t_m}(x))(D\eta_{s,t_m}(x)(h_{i_1}^{(q)})) d\beta_k(t_m) \dots d\beta_k(t_1), \\
& z_{s,t}^{m,N}(x)(h_{i_1}^{(q)}) \\
&= h_{i_1}^{(q)} + \sum_k \int_s^t \tilde{D}\tilde{A}_k(t_1, z_{s,t_1}^{N-1}(x))(h_{i_1}^{(q)}) + \sum_k \int_s^{t_1} \tilde{D}\tilde{A}_k(t_2, z_{s,t_2}^{N-2}(x))(h_{i_1}^{(q)}) \\
&\quad + \dots + \sum_k \int_s^{t_{m-1}} \tilde{D}\tilde{A}_k(t_m, z_{s,t_m}^{N-m}(x))(D\eta_{s,t_m}(x)(h_{i_1}^{(q)})) d\beta_k(t_m) \dots d\beta_k(t_1)
\end{aligned}$$

and taking  $N \geq m + n(\epsilon')$ , we have by Lemma 3, (3.10) and (4.9),

$$\begin{aligned}
& E[\|D\eta_{s,t}(x)(h_{i_1}^{(q)}) - Dz_{s,t}^N(x)(h_{i_1}^{(q)})\|_{-q}^{2j}] \\
&\leq 2^{2j-1} E[\|D\eta_{s,t}(x)(h_{i_1}^{(q)}) - y_{s,t}^{1,N}(x)(h_{i_1}^{(q)})\|_{-q}^{2j}] \\
&\quad + (2^{2j-1})^2 E[\|y_{s,t}^{1,N}(x)(h_{i_1}^{(q)}) - z_{s,t}^{1,N}(x)(h_{i_1}^{(q)})\|_{-q}^{2j}] \\
&\quad + \sum_{k=1}^{m-1} \{ (2^{2j-1})^{k+2} E[\|z_{s,t}^{k,N}(x)(h_{i_1}^{(q)}) - y_{s,t}^{k+1,N}(x)(h_{i_1}^{(q)})\|_{-q}^{2j}] \\
&\quad + (2^{2j-1})^{k+3} E[\|y_{s,t}^{k+1,N}(x)(h_{i_1}^{(q)}) - z_{s,t}^{k+1,N}(x)(h_{i_1}^{(q)})\|_{-q}^{2j}] \} \\
&\quad + (2^{2j-1})^{m+2} E[\|z_{s,t}^{m,N}(x)(h_{i_1}^{(q)}) - Dz_{s,t}^N(x)(h_{i_1}^{(q)})\|_{-q}^{2j}] \\
&\leq C_{43}(h_{i_1}^{(q)}) \{ \delta^{2j} (2^{2j-1})^2 RT + \epsilon' (2^{2j-1})^3 M^{2j} RT \\
&\quad + \sum_{k=1}^{m-1} ((2^{2j-1})^{2k+3} \delta^{2j} R^{k+1} M^{2jk}
\end{aligned}$$



$$\begin{aligned}
& + (2^{2j-1})^{2k+4} \epsilon' R^{k+1} M^{2j(k+1)} T^{k+1} / (k+1)! \\
& + (2^{2j-1})^{2m+2} R^m M^{2jm} T^m / m! \} \\
& \leq C_{44} (\delta^{2j} + \epsilon' + (2^{2j-1})^{2m+2} R^m M^{2jm} T^m / m!).
\end{aligned}$$

which gives (4.7) for  $k=1$  if we take sufficiently small  $\delta$ ,  $\epsilon'$  and large  $m$ . We assume (4.7) holds for integers  $1 \leq k \leq \ell$ ,  $\ell \geq 1$ .

Since

$$\begin{aligned}
& D^{\ell+1} (A_k(r, \eta_{s,r}(x))) (h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_{\ell+1}}^{(q)}) \\
& = D A_k(r, \eta_{s,r}(x)) (D^{\ell+1} \eta_{s,r}(x) (h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_{\ell+1}}^{(q)}))
\end{aligned}$$

+ finite sum of terms of the type

$$D^u A_k(r, \eta_{s,r}(x)) (D^{n_1} \eta_{s,r}(x) (h_{j_1(1)}^{(q)}, h_{j_2(1)}^{(q)}, \dots, h_{j_{n_1}(1)}^{(q)})),$$

$$D^{n_2} \eta_{s,r}(x) (h_{j_1(2)}^{(q)}, h_{j_2(2)}^{(q)}, \dots, h_{j_{n_2}(2)}^{(q)}), \dots,$$

$$D^{n_u} \eta_{s,r}(x) (h_{j_1(u)}^{(q)}, h_{j_2(u)}^{(q)}, \dots, h_{j_{n_u}(u)}^{(q)})).$$

where

$$2 \leq u \leq \ell+1, n_1 + n_2 + \dots + n_u = \ell+1, \{h_{j_{n_1}(1)}^{(q)}, i=1, 2, \dots, u\} = \{h_{i_j}^{(q)}, j=1, 2, \dots, \ell+1\}$$

and

$$D^{\ell+1} \eta_{s,t}(x) (h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_{\ell+1}}^{(q)})$$

$$= \sum_k \int_s^t D^{\ell+1}(A_k(r, \eta_{s,r}(x))) (h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_{\ell+1}}^{(q)}) d\beta_k(r).$$

so (4.7) for  $k \geq 2$  can be proved similarly.

Since  $F \in \mathfrak{A}_E$ , for any  $0 < \epsilon' < 1$ , we have a weighted Schwartz functional  $\tilde{F}(x) = f(\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_m \rangle)$  such that

$$(4.12) \quad \sum_{k=0}^{n+1} \sup_{x \in E'_q} e^{-\|x\|_{-q'}} \|D^k(F(x) - \tilde{F}(x))\|_{H.S.}^{(q')} < \epsilon'.$$

Then to prove Proposition 2, it is enough to show  $(U(t,s)F)(x)$  is approximated by weighted Schwartz functionals in  $\|\cdot\|_{p,k}^{(q)}$ ,  $0 \leq k \leq n$ . Since  $D^k(F(\eta_{s,t}(x))) (h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)})$  is a finite sum of terms of the type

$$(4.13) \quad \begin{aligned} & I_{h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)}}(\eta_{s,t}(x)) \\ &= D^u F(\eta_{s,t}(x)) (D^{n_1} \eta_{s,t}(x) (h_{j_1(1)}^{(q)}, h_{j_2(1)}^{(q)}, \dots, h_{j_{n_1}(1)}^{(q)}), D^{n_2} \eta_{s,t}(x) \\ & \quad (h_{j_1(2)}^{(q)}, h_{j_2(2)}^{(q)}, \dots, h_{j_{n_2}(2)}^{(q)}), \dots, D^{n_u} \eta_{s,t}(x) (h_{j_1(u)}^{(q)}, h_{j_2(u)}^{(q)}, \dots, h_{j_{n_u}(u)}^{(q)})). \end{aligned}$$

where  $0 \leq u \leq k$  and  $n_1 + n_2 + \dots + n_u = k$ , so that setting

$$(4.14) \quad \begin{aligned} & J_{h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)}}(z_{s,t}^N(x)) \\ &= D^u \tilde{F}(z_{s,t}^N(x)) (D^{n_1} z_{s,t}^N(x) (h_{j_1(1)}^{(q)}, h_{j_2(1)}^{(q)}, \dots, h_{j_{n_1}(1)}^{(q)}), D^{n_2} z_{s,t}^N(x) \\ & \quad (h_{j_1(2)}^{(q)}, h_{j_2(2)}^{(q)}, \dots, h_{j_{n_2}(2)}^{(q)}), \dots, D^{n_u} z_{s,t}^N(x) (h_{j_1(u)}^{(q)}, h_{j_2(u)}^{(q)}, \dots, h_{j_{n_u}(u)}^{(q)})). \end{aligned}$$

we see that  $(\|U(t,s)F - E[\tilde{F}(z_{s,t}^N(\cdot))]\|_{p,k}^{(q)})^2$  is dominated by a finite sum of terms of the type

$$\begin{aligned}
 (4.15) \quad & C_{45} \sup_{x \in E'_p} e^{-2\|x\|} \sum_{i_1, i_2, \dots, i_k=1}^{\infty} E[|I_{h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)}}(\eta_{s,t}(x)) \\
 & - J_{h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)}}(z_{s,t}^N(x))|^2] \\
 & \leq C_{46} \left( \sup_{x \in E'_p} e^{-2\|x\|} \sum_{i_1, i_2, \dots, i_k=1}^{\infty} E[e^{2\|\eta_{s,t}(x)\|} (\epsilon')^2 \|D^{n_1} \eta_{s,t}(x)(h_{j_1^{(1)}}^{(q)} \right. \\
 & \quad h_{j_2^{(1)}}^{(q)}, \dots, h_{j_{n_1}^{(1)}}^{(q)}\|_{-q}^2 \|D^{n_2} \eta_{s,t}(x)(h_{j_1^{(2)}}^{(q)}, h_{j_2^{(2)}}^{(q)}, \dots, h_{j_{n_2}^{(2)}}^{(q)}\|_{-q}^2 \\
 & \quad \dots \|D^{n_u} \eta_{s,t}(x)(h_{j_1^{(u)}}^{(q)}, h_{j_2^{(u)}}^{(q)}, \dots, h_{j_{n_u}^{(u)}}^{(q)}\|_{-q}^2) \\
 & \quad + \sup_{x \in E'_p} e^{-2\|x\|} \sum_{i_1, i_2, \dots, i_k=1}^{\infty} E[|J_{h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)}}(\eta_{s,t}(x)) \\
 & \quad \left. - J_{h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)}}(z_{s,t}^N(x))|^2] \right).
 \end{aligned}$$

Lemmas 5 and 6 and (4.12) give

$$\begin{aligned}
 (4.16) \quad & \sup_{x \in E'_p} e^{-\|x\|} \max\{E[(\|D^u \tilde{F}(z_{s,t}^N(x))\|_{H.S.}^{(q')})^2]^{1/2}, E[(\|D^{u+1} \tilde{F}(z_{s,t}^N(x)) \\
 & + \tau(\eta_{s,t}(x) - z_{s,t}^N(x))\|_{H.S.}^{(q')})^2]^{1/2}\} \leq C_{47}(T), \quad 0 \leq \tau \leq 1, \quad 0 \leq s, t \leq T.
 \end{aligned}$$

Hence from (3.10), (4.3), (4.16) and Lemma 5, we have constants  $C_{48}$  and  $C_{49}$  independent of  $\epsilon'$ , and for any  $\epsilon > 0$ , a natural number  $N_0$  such that (4.15) is

dominated by

$$\begin{aligned}
 (4.17) \quad & \epsilon/3 + C_{48}\epsilon' + C_{49} \sum_{i_1, i_2, \dots, i_k=1}^{N_0} E[\|\eta_{s,t}(x) - z_{s,t}^N(x)\|_{-q}^2 \cdot \|D^{n_1} \eta_{s,t}(x)(h_{j_1}^{(q)}(x)) \\
 & \quad h_{j_2}^{(q)}(x), \dots, h_{j_{n_1}}^{(q)}(x)\|_{-q}^2 \cdot \dots \|D^{n_u} \eta_{s,t}(x)(h_{j_1}^{(q)}(x), h_{j_2}^{(q)}(x), \dots, h_{j_{n_u}}^{(q)}(x))\|_{-q}^2 \\
 & \quad + \sum_{r=1}^u \|D^{n_1} z_{s,t}^N(x)(h_{j_1}^{(q)}(x), h_{j_2}^{(q)}(x), \dots, h_{j_{n_1}}^{(q)}(x))\|_{-q}^2 \cdot \dots \|D^{n_{r-1}} z_{s,t}^N(x)(h_{j_1}^{(q)}(x), \\
 & \quad h_{j_2}^{(q)}(x), \dots, h_{j_{n_{r-1}}}^{(q)}(x))\|_{-q}^2 \cdot \|D^{n_r} \eta_{s,t}(x)(h_{j_1}^{(q)}(x), h_{j_2}^{(q)}(x), \dots, h_{j_{n_r}}^{(q)}(x))\|_{-q}^2 \\
 & \quad - \|D^{n_r} z_{s,t}^N(x)(h_{j_1}^{(q)}(x), h_{j_2}^{(q)}(x), \dots, h_{j_{n_r}}^{(q)}(x))\|_{-q}^2 \cdot \|D^{n_{r+1}} \eta_{s,t}(x)(h_{j_1}^{(q)}(x), \\
 & \quad h_{j_2}^{(q)}(x), \dots, h_{j_{n_{r+1}}}^{(q)}(x))\|_{-q}^2 \cdot \dots \|D^{n_u} \eta_{s,t}(x)(h_{j_1}^{(q)}(x), h_{j_2}^{(q)}(x), \dots, h_{j_{n_u}}^{(q)}(x))\|_{-q}^2].
 \end{aligned}$$

Therefore noting (3.10), (4.6), (4.7), (4.9), (4.15), and (4.17) and taking sufficiently small  $\epsilon'$ ,  $\delta$  and large  $N$ , we obtain

$$\sup_{x \in E'_p} e^{-\|x\|} {}^{-p} \mathbb{D}^k((U(t,s)F)(x)) - D^k(E[\tilde{F}(z_{s,t}^N(x))]) \|_{H.S.}^{(q)} < \epsilon.$$

The rest is to prove that  $E[\tilde{F}(z_{s,t}^N(x))]$  is a weighted Schwartz functional. Of course  $E[\tilde{F}(z_{s,t}^N(x))] = \phi_{s,t}(\langle x, f_1 \rangle, \langle x, f_2 \rangle, \dots, \langle x, f_m \rangle, \langle x, f_1 \rangle, \langle x, f_2 \rangle, \dots, \langle x, f_\ell \rangle)$  is a smooth functional. To prove  $g(x)\phi_{s,t}(x) \in \mathcal{Y}(\mathbb{R}^{\ell+m})$ , by the Leibniz formula, it is sufficient to examine the finiteness of

$$\sup_{x \in \mathbb{R}^{\ell+m}} (1+|x|^2)^n |(\frac{d}{dx})^r g(x)(\frac{d}{dx})^k \phi_{s,t}(x)|, \text{ for any integers } 0 \leq r, k \leq n.$$

By the expression (4.14) of  $D^k(\tilde{F}(z_{s,t}^N(x)))(h_{i_1}^{(q)}, h_{i_2}^{(q)}, \dots, h_{i_k}^{(q)})$ , (4.9) and the fact that  $f(x) = h(x)\varphi(x)$ ,  $x \in \mathbb{R}^m$  and  $|(\frac{d}{dx})^r g(x)| \leq C_{50} \exp(-\sum_{i=1}^{\ell+m} \sqrt{|x_i|})$ , it is enough to show the finiteness of

$$(4.18) \quad \sup_Q (1 + \sum_{i=1}^m \langle x, \xi_i \rangle^2 + \sum_{j=1}^{\ell} \langle x, \zeta_j \rangle^2)^n \exp(-\sum_{i=1}^{\ell} \sqrt{|\langle x, \xi_i \rangle|} - \sum_{j=1}^m \sqrt{|\langle x, \zeta_j \rangle|}) \\ \times E[(h^{(\mu)}(z_{s,t}^N(x)) \varphi^{(v)}(z_{s,t}^N(x)))^2]^{1/2},$$

where

$$Q = \{x: (\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_m \rangle, \langle x, \zeta_1 \rangle, \langle x, \zeta_2 \rangle, \dots, \langle x, \zeta_{\ell} \rangle) \in \mathbb{R}^{\ell+m}\},$$

$$h^{(\mu)}(x) = (\frac{d}{dx})^{\mu} h(x), \quad \varphi^{(v)}(x) = (\frac{d}{dx})^v \varphi(x), \quad x \in \mathbb{R}^m,$$

$$\bar{h}^{(\mu)}(z_{s,t}^N(x)) = h^{(\mu)}(\langle z_{s,t}^N(x), \xi_1 \rangle, \langle z_{s,t}^N(x), \xi_2 \rangle, \dots, \langle z_{s,t}^N(x), \xi_m \rangle)$$

and

$$\bar{\varphi}^{(v)}(z_{s,t}^N(x)) = \varphi^{(v)}(\langle z_{s,t}^N(x), \xi_1 \rangle, \langle z_{s,t}^N(x), \xi_2 \rangle, \dots, \langle z_{s,t}^N(x), \xi_m \rangle).$$

Since  $|h^{(\mu)}(x)| \leq C_{51} \exp(-\sum_{i=1}^m \sqrt{|x_i|})$ , (4.11) of Lemma 6 yields that (4.18)

is dominated by

$$(4.19) \quad \sup_Q (1 + \sum_{i=1}^m \langle x, \xi_i \rangle^2 + \sum_{j=1}^{\ell} \langle x, \zeta_j \rangle^2)^n \exp(-\sum_{j=1}^{\ell} \sqrt{|\langle x, \zeta_j \rangle|}) E[(\bar{\varphi}^{(v)}(z_{s,t}^N(x)))^4]^{1/4} \\ \leq C_{52} \sup_Q (1 + \sum_{i=1}^m \langle x, \xi_i \rangle^2 + \sum_{j=1}^{\ell} \langle x, \zeta_j \rangle^2)^n \exp(-\sum_{j=1}^{\ell} \sqrt{|\langle x, \zeta_j \rangle|}) \\ \times E \left[ \frac{(1 + \sum_{i=1}^m \langle z_{s,t}^N(x), \xi_i \rangle^2)^{4n}}{(1 + \sum_{i=1}^m \langle z_{s,t}^N(x), \xi_i \rangle^2)^{4n}} |\bar{\varphi}^{(v)}(z_{s,t}^N(x))|^4 \right]^{1/4} \\ \leq C_{53} \|\varphi\|_n \sup_Q (1 + \sum_{i=1}^m \langle x, \xi_i \rangle^2 + \sum_{j=1}^{\ell} \langle x, \zeta_j \rangle^2)^n \exp(-\sum_{j=1}^{\ell} \sqrt{|\langle x, \zeta_j \rangle|})$$

$$\times E \left[ \frac{1}{(1 + \sum_{i=1}^m \langle z_{s,t}^N(x), F_i \rangle^2)^{4n}} \right]^{1/4}.$$

where  $\|\varphi\|_n = \sup_{\substack{x \in \mathbb{R}^m \\ 0 \leq r \leq n}} (1 + |x|^2)^n |\varphi^{(r)}(x)|$ .

On the other hand, we can verify the following lemma.

**Lemma 7.** For any  $F_1, F_2, \dots, F_m \in E$  and any integer  $p \geq 1$ , we have

$$E \left[ \frac{1}{(1 + \sum_{i=1}^m \langle z_{s,t}^N(x), F_i \rangle^2)^p} \right] \leq C_{54}(T) \frac{1}{(1 + \sum_{i=1}^m \langle x, F_i \rangle^2)^p}, \quad 0 \leq s, t \leq T.$$

**Proof.** Setting  $\theta(x) = \frac{1}{(1 + \sum_{i=1}^m \langle x, F_i \rangle^2)^p}$  and applying the Itô formula for

$\theta(z_{s,t}^N(x))$ , we get

$$\begin{aligned} (4.20) \quad E \left[ \frac{1}{(1 + \sum_{i=1}^m \langle z_{s,t}^N(x), F_i \rangle^2)^p} \right] &= \frac{1}{(1 + \sum_{i=1}^m \langle x, F_i \rangle^2)^p} \\ &+ E \left[ \int_s^t -2p(1 + \sum_{i=1}^m \langle z_{s,r}^N(x), F_i \rangle^2)^{-(p+1)} \left( \sum_{i=1}^m \langle z_{s,r}^N(x), F_i \rangle \langle \tilde{B}(r, z_{s,r}^{N-1}(x)), F_i \rangle \right) dr \right] \\ &+ E \left[ \int_s^t \sum_{j=1}^\infty (2p(p+1)(1 + \sum_{i=1}^m \langle z_{s,r}^N(x), F_i \rangle^2)^{-(p+2)} \left( \sum_{i=1}^m \langle z_{s,r}^N(x), F_i \rangle \langle \tilde{A}(r, z_{s,r}^{N-1}(x)) h_j^{(0)}, F_i \rangle \right)^2 \right. \right. \\ &\quad \left. \left. - p(1 + \sum_{i=1}^m \langle z_{s,r}^N(x), F_i \rangle^2)^{-(p+1)} \left( \sum_{i=1}^m \langle \tilde{A}(r, z_{s,r}^{N-1}(x)) h_j^{(0)}, F_i \rangle^2 \right) \right) dr \right]. \end{aligned}$$

By the boundedness of  $\tilde{A}_k(t, x)$ , (4.20) is dominated by

$$\frac{1}{(1 + \sum_{i=1}^m \langle z_{s,r}^N(x), F_i \rangle^2)^p} + C_{55} \int_s^t E \left[ \frac{1}{(1 + \sum_{i=1}^m \langle z_{s,r}^N(x), F_i \rangle^2)^p} \right] dr.$$

which yields the proof of the lemma, together with the Gronwall lemma.

Using this lemma, we see that the right hand side of (4.18) is dominated by

$$C_{56} \|\varphi\|_n \sup_Q (1 + \sum_{i=1}^m \langle x, \xi_i \rangle^2 + \sum_{j=1}^{\ell} \langle x, \zeta_j \rangle^2)^n \exp(-\sum_{j=1}^{\ell} \sqrt{|\langle x, \zeta_j \rangle|}) \\ \times \frac{1}{(1 + \sum_{i=1}^m \langle x, \xi_i \rangle^2)^n} < \infty.$$

Hence  $E[\tilde{F}(z_{s,t}^N(x))]$  is a weighted Schwartz functional and the proof of Proposition 2 is complete.

The following remark is immediate.

Remark. Under the assumptions of Proposition 2,  $(L(t)F)(x) \in \mathcal{D}_E$ , if  $F \in \mathcal{D}_E$ .

##### §5. Theorem

Propositions 1 and 2, together with Remark 1, yield

Theorem. Suppose that the coefficients  $A(t,x)$  and  $B(t,x)$  satisfy the conditions (H1)-(H3) and are approximated by bounded smooth functionals on  $E'$ . Then  $L(t)$  generates the Kolmogorov evolution operator  $U(t,s)$  from  $\mathcal{D}_E$  into itself. Further under the same assumption on the initial value as in Proposition 1, the continuous  $\mathcal{D}_E$ -process solution of (1.1) is uniquely given by

$$X_F(t) = X_{U(t,0)F}(0) + W_F(t) + \int_0^t W_L(s)U(t,s)F(s)ds.$$

As a direct application of our theorem, we give below another approach to

the fluctuation problem in [4].

**Example. Lattice system of interacting diffusions**

First we begin to explain the system that Deuschel considered in [4]. Let  $\mathbb{Z}^d$  be the  $d$ -dimensional lattice,  $i = (i_1, i_2, \dots, i_d) \in \mathbb{Z}^d$  and  $\mathcal{Y} = \mathcal{Y}(\mathbb{Z}^d)$  the Schwartz space of rapidly decreasing sequences  $\xi = (\xi_i)$ , metrized by the countably many semi-norms:

$$\|\xi\|_p^2 = \sum_{i \in \mathbb{Z}^d} (1 + |i|)^{2p} |\xi_i|^2, \quad p = 0, 1, 2, \dots$$

The dual space  $\mathcal{Y}' = \mathcal{Y}'(\mathbb{Z}^d)$  of  $\mathcal{Y}$  is the collection of all slowly increasing sequences  $x = (x_i)$  such that for some integer  $p \geq 0$ ,

$$\|x\|_{-p}^2 = \sum_{i \in \mathbb{Z}^d} (1 + |i|)^{-2p} |x_i|^2 < \infty.$$

Let  $b_i(x)$ ,  $i \in \mathbb{Z}^d$ , be a real valued infinitely many times  $\mathcal{Y}'$ -Fréchet differentiable mapping on  $\mathcal{Y}'$  for every integer  $p \geq 0$  such that  $b_i(x) = \hat{b}(\theta_i x)$ , where  $\hat{b}(x)$  is a real valued mapping on  $\mathcal{Y}'$  and  $\theta_i x = (x_{j+i})$ .

(V1) We have some natural number  $p_0$  such that

$$\sum_{i \in \mathbb{Z}^d} (1 + |i|)^{-2p_0} \left( \sup_{x \in \mathcal{Y}'} |b_i(x)| \right)^2 < \infty.$$

(V2) For any integers  $n \geq 1$  and  $p \geq 0$ ,

$$\sum_{i \in \mathbb{Z}^d} (1 + |i|)^{-2p} \left( \sup_{x \in \mathcal{Y}'} \|D^n b_i(x)\|_{H.S.}^{(p)} \right)^2 < \infty.$$

(V3) For any integers  $p \geq p_0$ ,  $q \geq 0$  and  $n \geq 0$ , there exists a sequence of real valued bounded smooth functionals  $b_i^{(m)}(x)$  such that

$$\lim_{m \rightarrow \infty} \sup_{x \in \mathcal{Y}'} \|D^n b_i(x) - D^n b_i^{(m)}(x)\|_{H.S.}^{(q)} = 0.$$



Let  $x(t) = (x_i(t), i \in \mathbb{Z}^d)$  be an  $\mathcal{Y}'(\mathbb{Z}^d)$ -valued solution of the following equation:

$$(5.1) \quad x_i(t) = \sigma_i + B_i(t) + \int_0^t b_i(x(s)) ds,$$

$$b_i(x) = \hat{b}(\theta_i x), \quad \theta_i x = (x_{j+i}),$$

where  $(B_i(t))$  are independent copies of the 1-dimensional standard Brownian motion  $B(t)$ ,  $(\sigma_i)$  are independent copies of the 1-dimensional random variable  $\sigma$  independent of  $B(t)$  and for any  $\epsilon > 0$ ,  $E[\exp(\epsilon \|\sigma_i\|_{-p_0})] < \infty$ . For a finite lattice  $V \in \mathbb{Z}^d$ , consider

$$T_V(t) = |V|^{-1/2} \sum_{i \in V} \delta_{\theta_i x(t)}.$$

Now put

$$\langle U_V(t), \phi \rangle = \langle T_V(t), \phi \rangle - E[\langle T_V(t), \phi \rangle], \quad \phi \in C_0^\infty(\mathcal{Y}').$$

Then it can be proved (see [18], [22]) that  $U_V(t)$  becomes a strongly continuous  $C_0^\infty(\mathcal{Y}')$ -valued stochastic process. We will prove tightness for  $U_V(t)$ ,  $V \in \mathbb{Z}^d$  following [5], [19], in  $C([0, \infty); C_0^\infty(\mathcal{Y}'))$ . Let  $L_0$  be an operator defined by

$$(L_0 F)(x) = \frac{1}{2} \text{trace}_{\ell^2(\mathbb{Z}^d)} D^2 F(x) + DF(x)(b(x)), \quad F \in \mathfrak{D}_{\mathcal{Y}'(\mathbb{Z}^d)},$$

where  $b(x) = (b_i(x))$ .

By the conditions (V1) and (V2), equation (5.1) is solved in  $\mathcal{Y}'_{p_0}$ , so that  $x(t) \in \mathcal{Y}'_{p_0}$ . Then we have by the exponential integrability proved in the same way as in Lemma 5,

$$E[\langle T_V(t), \phi \rangle^2] \leq C_{57} \|\phi\|_{p_0, 0, 0}^2.$$

Since  $C_0^\infty(\mathcal{Y}')$  is dense in  $\mathfrak{D}_{\mathcal{Y}'}(\mathbb{Z}^d)$ ,  $T_V(t)$  is extended to a continuous  $\mathcal{L}(\mathfrak{D}_{\mathcal{Y}'}(\mathbb{Z}^d))$ -process. We denote the extension by  $T_{\phi,V}(t)$ .

Let  $\phi(x) = \phi(\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_n \rangle)$ ,  $\phi \in S(\mathbb{R}^n)$ . By the Itô formula, we get

$$(5.2) \quad \langle T_V(t), \phi \rangle - \langle T_V(0), \phi \rangle = M_{\phi,V}(t) + \int_0^t T_{L_0} \phi, V(s) ds,$$

where

$$M_{\phi,V}(t) = |V|^{-1/2} \sum_{i \in V} \int_0^t \sum_{j=1}^n \frac{\partial}{\partial x_j} \phi(\langle \theta_i x(s), \xi_1 \rangle, \langle \theta_i x(s), \xi_2 \rangle, \dots, \langle \theta_i x(s), \xi_n \rangle) \\ \left( \sum_{k \in \mathbb{Z}^d} \xi_k^1 dB(s) \right),$$

where  $\xi_i = (\xi_k^1)$ ,  $k \in \mathbb{Z}^d$ .

From the independence of  $B_i(t)$ ,  $i \in V$  and the fact that  $x(t) \in \mathcal{Y}'_{p_0}$ , we have for  $t \in [0, T]$ ,

$$(5.3) \quad E[M_{\phi,V}(t)^4] \leq C_{58} \|\phi\|_{p_0,1,1}^4.$$

Then  $M_{\phi,V}(t)$  can be extended to a continuous  $\mathcal{L}(\mathfrak{D}_{\mathcal{Y}'}(\mathbb{Z}^d))$ -process and has the same regularity properties that the  $\mathcal{L}(\mathfrak{D}_{\mathcal{Y}'}(\mathbb{Z}^d))$ -Wiener process has. Conditions (V1)-(V3) guarantee that  $L_0$  belongs to the class dealt with in the Theorem. We use the same notation  $U(t,s)$  to represent the evolution operator generated by  $L_0$ . Thus the solution of (5.2) is given as follows:

$$\langle T_V(t), \phi \rangle = T_{U(t,0)} \phi, V(0) + M_{\phi,V}(t) + \int_0^t T_{L_0} U(t,s) \phi, V(s) ds$$

as in the proof of Proposition 1. Hence by (5.3) and the Kolmogorov test for a real Wiener process, we get

$$E[|\langle U_V(t) - U_V(s), \phi \rangle|^4] \leq C_{59} |t-s|^2$$

and further

$$E[|\langle U_V(t), \phi \rangle|^2] \leq C_{60} \{ \|\phi\|_{p_0, 1, 1}^2 + \sup_{0 \leq s \leq t} \|U(t, s)\phi\|_{p_0, 1, 3}^2 \}$$

which proves the tightness in  $C([0, \infty); C_0^\infty(\mathcal{Y}'))'$ . ([5], [18]). By the Skorokhod theorem and the usual limiting argument, the limit process  $N(t)$  of  $U_V(t)$  satisfies the SDE

$$(5.4) \quad \langle N(t) - N(0), \phi \rangle = W_\phi(t) + \int_0^t N_{L_0} \phi(s) ds,$$

where  $N_F(t)$ ,  $F \in \mathcal{D}_{\mathcal{Y}'(\mathbb{Z}^d)}$ , is the extension of  $N(t)$  and  $W_F(t)$  is a  $\mathcal{L}(\mathcal{D}_{\mathcal{Y}'(\mathbb{Z}^d)})$ -Wiener process [8].

The uniqueness for solutions of the equation (5.4) discussed in Theorem implies the identification of the distribution of the limit process, ([20], [21]), which implies that  $U_V(t)$  converges to a Gaussian field in  $C([0, \infty); C_0^\infty(\mathcal{Y}'))'$ .

56. A fluctuation theorem for a system of interacting, spatially distributed neurons.

A problem in neurophysiology that has received considerable attention in recent years, is the stochastic behavior of the voltage potential of a spatially distributed neuron [11, 26]. When the spatial dimension of the neuronal membrane is greater than one, the voltage potential is modeled as a stochastic process taking values in the dual of some nuclear space such as the space of Schwartz distributions  $\mathcal{Y}'(\mathbb{R}^d)$ . The SDE satisfied by the voltage potential is best introduced via the following general model: Let  $H$  be a real separable Hilbert space, in applications, usually  $H = L^2(\mathfrak{X}, d\mu)$  where  $\mathfrak{X}$  is the membrane of the spatially extended neuron (e.g.  $\mathfrak{X} = [0, b]$ , a  $d$ -dimensional rectangle or a compact Riemannian manifold with or without boundary, and  $\mu$  is

the appropriate natural measure on  $\mathfrak{A}$ ). Let  $T_t$  be a strongly continuous semigroup on  $H$  generated by a closed, densely defined operator  $\mathfrak{A}$  such that  $(\mathfrak{A}\xi, \xi)_H \leq 0$  for  $\xi \in \text{Dom}(\mathfrak{A})$  where  $(\cdot, \cdot)_H$  denotes the inner product of  $H$ . Assume that some power of the resolvent of  $\mathfrak{A}$  is a Hilbert-Schmidt operator i.e.

$$(6.1) \quad (\lambda I - \mathfrak{A})^{-r_1} \text{ is Hilbert-Schmidt for some } r_1 > 0.$$

Then there is a CONS  $\{\varphi_j\}_{j \geq 1}$  in  $H$  such that  $-\mathfrak{A}\varphi_j = \lambda_j \varphi_j$  for any  $j \geq 1$  and  $0 \leq \lambda_1 \leq \lambda_2 \leq \dots$ . Set

$$E = \{\xi \in H; \sum_{j=1}^{\infty} (1+\lambda_j)^{2r} (\xi, \varphi_j)_H^2 < \infty \text{ for any } r \geq 0\}.$$

Define the inner product on  $E$ ,

$$(\xi, \zeta)_r = \sum_{j=1}^{\infty} (1+\lambda_j)^{2r} (\xi, \varphi_j)_H (\zeta, \varphi_j)_H$$

and  $E_r$  as the  $\|\cdot\|_r$ -completion of  $E$ . ( $\|\xi\|_r^2 = (\xi, \xi)_r$ ) and  $E'_r$  as the dual of the Hilbert space  $E_r$ . For  $r < s$ ,  $E_s \subseteq E_r$  and  $E_0 = H$ . Condition (6.1) implies that the canonical injection  $E_p \rightarrow E_r$  is Hilbert-Schmidt if  $p > r + r_1$ . Hence  $E$  is nuclear.

Since  $\mathfrak{A}$  generates  $T_t$  on  $H$ , we have for  $\xi \in E$  and  $t > 0$ ,

$$T_t \xi = \sum_{j=1}^{\infty} e^{-t\lambda_j} (\xi, \varphi_j)_H \varphi_j.$$

The following properties of  $T_t$  can be easily verified:

- (a)  $T_t E \subseteq E$ ;
- (b) The restriction of  $T_t$  to  $E$  is an  $E$ -continuous semigroup;
- (c)  $t \rightarrow T_t \xi$  is continuous for every  $\xi \in E$ ;
- (d) The restriction of  $\mathfrak{A}$  on  $E$  maps  $E$  into  $E$  and is the generator of the

semigroup  $T_t$  on  $E$ ;

(e) For any  $\xi \in E$  and  $t > 0$ ,

$$\|T_t \xi\|_r \leq \|\xi\|_r.$$

The voltage potential is then derived as the solution of an  $E'$ -valued SDE

$$(6.2) \quad dX(t) = d\beta(t) + \mathcal{A}'X(t)dt$$

where  $\mathcal{A}'$  is the adjoint of  $\mathcal{A}$  on  $E$  and  $\beta(t)$  is an  $E'$ -valued Wiener process with  $E[\langle \beta(t), \xi \rangle \langle \beta(s), \zeta \rangle] = (t \wedge s)Q(\xi, \zeta)$ ,  $Q$  being an  $E$ -continuous quadratic form.

Let us now define

$$\langle V(t)x, \xi \rangle = \langle x, T_t \xi \rangle \quad \forall x \in E', \xi \in E.$$

Then, using property (e) above we have

$$\|V(t)x\|_{-r} = \sup_{\|\xi\|_r \leq 1} |\langle x, T_t \xi \rangle| \leq \|x\|_{-r} \sup_{\|\xi\|_r \leq 1} \|T_t \xi\|_r \leq \|x\|_{-r}$$

and so

$$(6.3) \quad \sup_{0 \leq t \leq T} \|V(t)x\|_{-r} \leq \|x\|_{-r}.$$

(6.3) is a special case of condition (VI2) below, which is thus satisfied for the class of spatially extended neurons whose voltage potentials are modeled by (6.2). For specific examples of  $L^2(\mathcal{A}, d\mu)$  and the semigroup  $T_t$  which describes the deterministic part of the behavior of the neuron, see [11].

We now come to the question of interacting assemblies of a very large number of neurons. This appears to be a very important problem of physiological interest since such large systems are involved in the functioning of the central nervous system. The difficulty consists in discovering the precise nature of the interaction in a mathematical form. In this section we consider an interaction similar to the mean-field interaction in particle

diffusions. Another, possibly more realistic interaction known in the physiological literature as "parallel fiber interaction" will be investigated in our future work.

Let  $b(x,y)$  be a mapping from  $E' \times E'$  to some  $E'_{p_6}$  such that  $b(\cdot, \cdot)$  is infinitely many  $E'_p$ -Fréchet differentiable for every integer  $p \geq 0$  and with all derivatives bounded;

$$(VI1) \quad \sup_{x,y \in E'} \|D_x^k D_y^m b(x,y)\|_{H.S.}^{(p)} < \infty$$

for any integers  $k, m$  and  $p \geq 0$ . Here  $D_x$  and  $D_y$  denote the Fréchet derivatives with respect to variables  $x$  and  $y$ . The  $i$ -th component  $X_i^{(n)}(t)$  of the  $n$ -system of diffusions is obeyed by the following stochastic differential equation:

$$(6.4) \quad dX_i^{(n)}(t) = d\beta_i(t) + \{A(t)X_i^{(n)}(t) - \frac{1}{n} \sum_{j=1}^n b(X_i^{(n)}(t), X_j^{(n)}(t))\}dt, \\ i=1, 2, \dots, n,$$

where  $\{\beta_i(t)\}$  are independent copies of an  $E'$ -valued Wiener process  $\beta(t)$ .

Suppose that  $A(t)$  generates the strongly continuous contraction evolution operator  $V(t,s)$  from  $E'$  to itself such that for any integer  $p$  and any  $T > 0$ , there exists some integer  $n(p,T) \geq p$  satisfying

$$(VI2) \quad \sup_{0 \leq s \leq t \leq T} \|V(t,s)x\|_{-n(p,t)} \leq \|x\|_{-p}.$$

Without loss of generality, we may assume  $n(p,T) \leq n(q,T)$  if  $p \leq q$ . Then (6.4) is equivalent to

$$(6.5) \quad X_i^{(n)}(t) = V(t,0)\sigma_i + \int_0^t V(t,s)d\beta_i(s) + \int_0^t V(t,s) \left( \frac{1}{n} \sum_{j=1}^n b(X_i^{(n)}(s), X_j^{(n)}(s)) \right) ds.$$

We assume the initial value  $\sigma_i$  to be an independent copy of  $\sigma$  such that

$E[\exp(\epsilon \| \cdot \|_{-p_7})] < \infty$  for every  $\epsilon > 0$  and some natural number  $p_7$ .

Since  $Q(\xi, \xi)$  is a continuous quadratic form on  $E$ , there exists an integer  $r$  such that

$$Q(\xi, \xi) = (Q_r^{1/2} \xi, Q_r^{1/2} \xi)_r$$

where  $Q_r$  is a self-adjoint operator on  $E_r$ . Then clearly,  $\beta(t) \in E'_{p_8}$  for some integer  $p_8 > r$ .

The solution of (6.5) until time  $T$  is easily obtained by the usual method of successive approximations in  $E'_{n(p_9, T)}$ , where  $p_9 = p_6 \vee p_7 \vee p_8$ .

For the finite measure  $\nu(dx)$  on  $E'$ , set  $b[x, \nu] = \int_E b(x, y) \nu(dy)$ , where the integral is the Bochner integral on  $E'$  and consider

$$(6.6) \quad \begin{aligned} dX_1(t) &= d\beta_1(t) + \{X(t)X_1(t) + b[X_1(t), \nu]\}dt, \\ \nu(t, dx) &= \text{the distribution of } X_1(t). \end{aligned}$$

Then according to the following lemma the empirical distribution

$\frac{1}{n} \sum_{j=1}^n \delta_{X_j^{(n)}(t)}$  converges to  $\nu(t, dx)$  in probability in the usual weak convergence

of measures, where  $\delta_x$  is the Dirac measure at  $x$  in  $E'$ .

Lemma 8. For any  $T > 0$  and integer  $j \geq 1$ ,

$$E[\|X_1^{(m)}(t) - X_1(t)\|_{-n(p_6, T)}^{2j}] \leq C_{61}(T)/m^j, \quad 0 \leq t \leq T.$$

Proof. Put  $n_0 = n(p_6, T)$ . Then the condition (VII) yields

$$\begin{aligned} & \|b(X_1^{(m)}(t), X_j^{(m)}(t)) - b(X_1(t), X_j^{(m)}(t))\|_{-p_6} \\ & \leq \sup_{x, y \in E'} \|D_x b(x, y)\|_{H.S.}^{(n_0)} \|X_1^{(m)}(t) - X_1(t)\|_{-n_0} \end{aligned}$$

$$\leq C_{62} \|X_1^{(m)}(t) - X_1(t)\|_{-n_0}$$

and

$$\|b(X_1(t), X_j^{(m)}(t)) - b(X_1(t), X_j(t))\|_{-p_6} \leq C_{63} \|X_j^{(m)}(t) - X_j(t)\|_{-n_0},$$

so that we have

$$\begin{aligned} (6.7) \quad & E[\|X_1^{(m)}(t) - X_1(t)\|_{-n_0}^{2j}] \\ & \leq C_{64}(T) \int_0^t E[\|V(t, s)\|_{-p_6}^{2j} \left\{ \frac{1}{m} \sum_{j=1}^m b(X_1^{(m)}(s), X_j^{(m)}(s)) - b[X_1(s), u] \right\} \|_{-n_0}^{2j} \}] ds \\ & \leq C_{65}(T) \int_0^t [E[\|X_1^{(m)}(s) - X_1(s)\|_{-n_0}^{2j}] + \frac{1}{m} \sum_{j=1}^m E[\|X_j^{(m)}(s) - X_j(s)\|_{-n_0}^{2j}]] \\ & \quad + E[\| \frac{1}{m} \sum_{j=1}^m \{b(X_1(s), X_j(s)) - b[X_1(s), u]\} \|_{-p_6}^{2j}] ds. \end{aligned}$$

Noticing the independence of  $X_i(t)$ ,  $i=1, 2, \dots, m$  and the condition (VI1), we have

$$(6.8) \quad E[\| \frac{1}{m} \sum_{j=1}^m \{b(X_1(s), X_j(s)) - b[X_1(s), u]\} \|_{-p_6}^{2j}] \leq C_{66}(T)/m^j.$$

Therefore Gronwall's inequality, together with (6.7) and (6.8), implies the assertion of Lemma 8.

Now we proceed to the discussion of the fluctuation problem. We are able to consider  $U_n(t) = \sqrt{n} \left( \frac{1}{n} \sum_{j=1}^n \delta_{X_j^{(n)}(t)} - u(t, dx) \right)$  as a  $C_0^\infty(E')'$ -valued continuous stochastic process [18], [22]. To check the tightness of  $U_n(t)$  in  $C([0, \infty); C_0^\infty(E')')$  of all continuous mappings from  $[0, \infty)$  into  $C_0^\infty(E')'$ , it is enough to verify the Kolmogorov tightness criterion for  $\langle U_n(t), \phi \rangle$ ,  $\phi \in C_0^\infty(E')$ , where  $\langle \cdot, \cdot \rangle$  denotes the canonical bilinear form on  $C_0^\infty(E')' \times C_0^\infty(E')$ . [5], [19].

We have the following exponential integrability.



Lemma 9. For any  $\alpha > 0$ ,  $T > 0$  and any integer  $p \geq n(p_0, T)$ , there exists a constant  $C_{67} = C_{67}(\alpha, T, p)$  such that

$$\sup_{0 \leq t \leq T} E[e^{\alpha \|X_1^{(n)}(t)\|^{-p}}] \vee E[e^{\alpha \|X_1(t)\|^{-p}}] \leq C_{67}.$$

Proof. Set  $n_0 = n(p_0, T)$ . Assumptions (VI1) and (VI2) give

$$\max\{\|X_1^{(n)}(t)\|_{-n_0}, \|X_1(t)\|_{-n_0}\} \leq \|\sigma_1\|_{-p_7} + C_{68} + \|\int_0^t V(t, s) dB_1(s)\|_{-n_0}$$

and hence the lemma can be proved in the same way as Lemma 5.

Once we know Lemmas 8 and 9, we can check the moment condition;

$$(6.9) \quad E[|\langle U_n(t) - U_n(s), \phi \rangle|^4] \leq C_{69}(\phi) |t-s|^2.$$

(see [8]). Similarly we have

$$(6.10) \quad \sup_{0 \leq t \leq T} E[\langle U_n(t), \phi \rangle^2] \leq C_{70}(T) \|\phi\|_{n(p_0, T), n(p_0, T), 1}^2.$$

Then a subsequence of  $U_n(t)$  converges to  $U(t)$  in  $C([0, \infty); C_0^\infty(E'))'$ . Further (6.9) and (6.10) guarantee that  $U_n(t)$  and  $U(t)$  can be extended to continuous  $\mathcal{L}(\mathfrak{H}_E)$ -processes and so we denote the extensions by  $(U_n)_F(t)$  and  $U_F(t)$ ,  $F \in \mathfrak{H}_E$ , respectively.

For any  $F \in \mathfrak{H}_E$ , define

$$\begin{aligned} (K(t)F)(x) = & \frac{1}{2} \text{trace}_{E_r} D^2 F(x) \circ [(Q_r^{1/2})^* x (Q_r^{1/2})^*] + DF(x)(b[x, u] + \mathfrak{A}(t)x) \\ & + \int_E DF(y)(b(y, x))u(t, dy) \end{aligned}$$

and

$$W_F(t) = U_F(t) - U_F(0) - \int_0^t U_{K(s)F}(s) ds.$$

where  $\text{trace}_{E_r} D^2 F(x) \circ [(Q_r^{1/2})^* x (Q_r^{1/2})^*] = \sum_{j=1}^{\infty} D^2 F(x) ((Q_r^{1/2})^* h_j(r), (Q_r^{1/2})^* h_j(r))$   
and  $*$  means the adjoint operator with respect to the dual pair on  $E' \times E$ .

By following the argument of [8] word by word, we have the proof that  $W_F(t)$  is a continuous  $\mathcal{V}(\mathfrak{H}_E)$ -Wiener process. Thus any limit process of convergent subsequences of  $U_n(t)$  satisfies the weak SDE of type (1.1).

Now we impose a rather technical condition on  $b(x, y)$ .

(VI3) For any  $\epsilon > 0$  and any integers  $p, q, n \geq 0$ , there exists a  $C_b^\infty$ -function  $\bar{b}(x, y)$  of  $\mathbb{R}^m \times \mathbb{R}^{m'}$  to  $E'_{p_6}$  such that

$$\sup_{x \in E'_p} \| D_x^\mu D_y^\nu [b(x, y) - \bar{b}(\langle x, \xi_1 \rangle, \langle x, \xi_2 \rangle, \dots, \langle x, \xi_m \rangle,$$

$$\langle y, \zeta_1 \rangle, \langle y, \zeta_2 \rangle, \dots, \langle y, \zeta_m \rangle)] \|_{H.S.}^{(q)} < \epsilon.$$

$$0 \leq \mu + \nu \leq n, \xi_i, \zeta_j \in E, i=1, 2, \dots, m \text{ and } j=1, 2, \dots, m'.$$

Here  $C_b^\infty$ -function means  $\bar{b}(x, y)$  itself and all the derivatives are bounded.

We set

$$(A(t)F)(x) = \frac{1}{2} \text{trace}_{E_r} D^2 F(x) \circ [(Q_r^{1/2})^* x (Q_r^{1/2})^*] + DF(x)(b[x, u] + \mathfrak{A}(t)x)$$

and

$$(J(t)F)(x) = \int_E DF(y)(b(y, x))u(t, dy).$$

Though  $\mathfrak{A}(t)x$  is not bounded, from a part of the proof of Proposition 2 and the assumptions (VI1) and (VI3), we can show

$$A(t)\mathfrak{H}_E \subset \mathfrak{H}_E, \quad \text{and} \quad J(t)\mathfrak{H}_E \subset \mathfrak{H}_E.$$

Since

$$\eta_{s,t}(x) = V(t, s)x + \int_s^t V(t, r)d\beta(r) + \int_s^t V(t, r)b[\eta_{s,r}(x), u]dr,$$

choosing  $q' > n(q'', T)$  such that  $q'' > \max\{p, p_0, q\}$  and  $\sum_{j=1}^{\infty} \|h_j^{(q)}\|_{-q''}^2 < \infty$  in the proofs of Propositions 1 and 2 and recalling the condition (VI2), we conclude

that Lemma 5 holds if  $q'$  in the right hand side is replaced by  $q''$  and hence, together with (VI1) and (VI3), we obtain that  $A(t)$  generates the Kolmogorov evolution operator from  $\mathfrak{D}_E$  into itself similarly. Further since  $J(t)$  satisfies the condition of Proposition 2 in [21] and the proof of Proposition 2 in [21] is valid for any Fréchet space,  $K(t) = A(t) + J(t)$  generates the Kolmogorov evolution operator  $U(t, s)$  from  $\mathfrak{D}_E$  into itself. Since the Theorem gives the identification of the distributions of the limit processes  $U(t)$ , we obtain the conclusion that under the assumptions (VI1) - (VI3) and the exponential integrability of  $\sigma$ ,  $U_n(t)$  converges to a Gaussian field governed by the weak SDE of type (1.1) in  $C([0, \infty); C_0^\infty(E'))'$ , namely,

$$dX_F(t) = dW_F(t) + X_{K(t)F}(t)dt,$$

where  $W_F(t)$  is an  $\mathcal{L}(\mathfrak{D}_E)$ -Wiener process with

$$E[W_F(t)W_G(s)] = \int_0^{t \wedge s} \left( \int_{E'} \sum_{j=1}^{\infty} DF(x) ((Q_r^{1/2})^* h_j(r)) DG(x) ((Q_r^{1/2})^* h_j(r)) u(\tau, dx) \right) d\tau$$

and  $X_F(0)$  is a Gaussian random variable with

$$E[X_F(0)X_G(0)] = E[F(\sigma)G(\sigma)] - E[F(\sigma)]E[G(\sigma)], \quad F, G \in \mathfrak{D}_E.$$

**Acknowledgements.** The authors wish to thank Professors D. Dawson and L. Gorostiza for valuable discussions of the problems studied in this paper. The second author would also like to express his appreciation to Professor T. Shiga for his many helpful suggestions.

## References

- [1] P. Billingsley: *Convergence of Probability Measures*, Wiley, New York-London-Sydney-Toronto, 1968.
- [2] T. Bojdecki and L.G. Gorostiza: Langevin equation for  $\mathcal{V}$ -valued Gaussian processes and fluctuation limits of infinite particle systems. *Probab. Th. Rel. Fields* 73 (1986), 227-244.
- [3] D.A. Dawson: Critical dynamics and fluctuations for a mean-field model of cooperative behavior. *J. Statist. Phys.* 31 (1983), 29-85.
- [4] J.D. Deuschel: Central limit theorem for an infinite lattice system of interacting diffusion processes. Preprint. 1987.
- [5] J.P. Fouque: La convergence et loi pour les processus a valeurs dans un espace nucleaire. *Ann. IHP* 20 (1984), 225-245.
- [6] I.M. Gelfand and G.E. Shilov: *Generalized functions 2*. Academic Press, New York and London, 1964.
- [7] J.I. Gikhman and A.V. Skorokhod: *Stochastic Differential Equations*, Springer, Berlin, 1972.
- [8] M. Hitsuda and I. Mitoma: Tightness problem and stochastic evolution equation arising from fluctuation phenomena for interacting diffusions. *J. Multivariate Anal.* (1986), 311-328.
- [9] R. Holley and D.W. Stroock: Central limit phenomena of various interacting systems. *Ann. Math.* 110 (1979), 333-393.
- [10] K. Ito: Infinite dimensional Ornstein-Uhlenbeck processes. Taniguchi Symp. SA, Katata, 197-224, Kinokuniya, Tokyo 1984.
- [11] G. Kallianpur and R. Wolpert: Infinite dimensional stochastic models for spatially distributed neurons. *Appl. Math. Optim.* 12 (1984) 125-172.
- [12] H. Komatsu: Semi-groups of operators in locally convex spaces. *J. Math. Soc. Japan*, 16 (1964), 232-262.
- [13] H. Kunita: Stochastic differential equations and stochastic flows of diffeomorphisms. *Lect. Notes in Math.* 1097, Springer 1984.
- [14] H.H. Kuo: Gaussian measures on Banach spaces. *Lect. Notes in Math.* 463, Springer, Berlin, 1975.
- [15] H.H. Kuo: Stochastic integrals in abstract Wiener space II. regularity properties. *Nagoya Math. J.* 50 (1973), 89-116.
- [16] H.P. McKean: Propagation of chaos for a class of non-linear parabolic equations. *Lecture series in Differential Equations* 7, Catholic Univ. (1967), 41-57.

- [17] R.A. Minlos: Generalized random processes and their extension to a measure. Selected Transl. Math. Statist. Probab. 3 (1962), 291-313.
- [18] I. Mitoma: On the sample continuity of  $\mathcal{G}'$ -processes. J. Math. Soc. Japan. (1983), 629-636.
- [19] I. Mitoma: Tightness of probabilities on  $C([0,1];\mathcal{G}')$  and  $D([0,1];\mathcal{G}')$ . Ann. Probab. 11 (1983), 989-999.
- [20] I. Mitoma: An  $\infty$ -dimensional inhomogeneous Langevin's equation. J. Functional Analysis 61 (1985), 342-359.
- [21] I. Mitoma: Generalized Ornstein-Uhlenbeck process having a characteristic operator with polynomial coefficients. Probab. Th. Rel. Fields 76 (1987), 533-555.
- [22] C. Martias: Sur les support des processus a valeurs dans des espaces nucleaires. To appear in Ann. IHP.
- [23] T. Shiga and H. Tanaka: Central limit theorem for a system of Markovian particles with mean-field interactions. Z. Wahrsch. verw. Gebiete 69 (1985), 439-459.
- [24] H.H. Schaefer: Topological vector spaces. Springer, Berlin 1971.
- [25] H. Totoki: A method of construction for measures on function spaces and its applications to stochastic processes. Mem. Fac. Sci. Kyushu Univ. Ser. A, Math. 15 (1962), 178-190.
- [26] J. Walsh: An introduction to stochastic partial differential equations. École d'Été de Probabilités de Saint-Flour XIV. Lect. Notes in Math. 1180. Springer, Berlin 1984.

206. E. Willekens, Estimation of convolution tails, Sept. 87.
207. J. Rosinski, On path properties of certain infinitely divisible processes, Sept. 87.
208. A.H. Korezlioglu, Computation of filters by sampling and quantization, Sept. 87.
209. J. Bather, Stopping rules and observed significance levels, Sept. 87.
210. S.T. Rachev and J.E. Yukich, Convolution metrics and rates of convergence in the central limit theorem, Sept. 87. *Ann. Probability*, to appear.
211. M. Fujisaki, Normed Bellman equation with degenerate diffusion coefficients and its applications to differential equations, Oct. 87.
212. G. Simons, Y.C. Yao and X. Wu, Sequential tests for the drift of a Wiener process with a smooth prior, and the heat equation, Oct. 87.
213. R.L. Smith, Extreme value theory for dependent sequences via the Stein-Chen method of Poisson approximation, Oct. 87.
214. C. Houdré, A vector bimeasure integral with some applications, June 88 (Revised).
215. M.R. Leadbetter, On the exceedance random measures for stationary processes, Nov. 87.
216. M. Marques, A study on Lebesgue decomposition of measures induced by stable processes, Nov. 87 (*Dissertation*).
217. M.T. Alpuim, High level exceedances in stationary sequences with extremal index, Dec. 87. *Stochastic Proc. Appl.*, to appear.
218. R.F. Serfozo, Poisson functionals of Markov processes and queueing networks, Dec. 87.
219. J. Bather, Stopping rules and ordered families of distributions, Dec. 87.
220. S. Cambanis and M. Maejima, Two classes of self-similar stable processes with stationary increments, Jan. 88.
221. H.P. Hücke, G. Kallianpur and R.L. Karandikar, Smoothness properties of the conditional expectation in finitely additive white noise filtering, Jan. 88. *J. Multivariate Anal.*, to appear.
222. I. Mitoma, Weak solution of the Langevin equation on a generalized functional space, Feb. 88.
223. L. de Haan, S.J. Resnick, H. Rootzén and C. de Vries, Extremal behaviour of solutions to a stochastic difference equation with applications to arch-processes, Feb. 88.
224. O. Kallenberg and J. Szulga, Multiple integration with respect to Poisson and Lévy processes, Feb. 88.
225. D.A. Dawson and L.G. Gorostiza, Generalized solutions of a class of nuclear space valued stochastic evolution equations, Feb. 88.
226. G. Samorodnitsky and J. Szulga, An asymptotic evaluation of the tail of a multiple symmetric  $\alpha$ -stable integral, Feb. 88.
227. J.J. Hunter, The computation of stationary distributions of Markov chains through perturbations, Mar. 88.
228. H.C. Ho and T.C. Sun, Limiting distribution of nonlinear vector functions of stationary Gaussian processes, Mar. 88.
229. R. Brigola, On functional estimates for ill-posed linear problems, Apr. 88.
230. M.R. Leadbetter and S. Nandagopalan, On exceedance point processes for stationary sequences under mild oscillation restrictions, Apr. 88.
231. S. Cambanis, J. P. Nolan and J. Rosinski, On the oscillation of infinitely divisible processes, Apr. 88.
232. G. Hardy, G. Kallianpur and S. Ramasubramanian, A nuclear space-valued stochastic differential equation driven by Poisson random measures, Apr. 88.
233. D.J. Daley, T. Rolski, Light traffic approximations in queues (II), May 88.
234. G. Kallianpur, I. Mitoma, R.L. Wolpert, Diffusion equations in duals of nuclear spaces, July 88.
235. S. Cambanis, Admissible translates of stable processes: A survey and some new models, July 88.
236. E. Platen, On a wide range exclusion process in random medium with local jump intensity, Aug. 88.
237. R.L. Smith, A counterexample concerning the extremal index, Aug. 88.
238. G. Kallianpur and I. Mitoma, A Langevin-type stochastic differential equation on a space of generalized functionals, Aug. 88.
239. C. Houdré, Harmonizability,  $\chi$  boundedness, (2,P)-boundedness of stochastic processes, Aug. 1988.
240. G. Kallianpur, Some remarks on Hu and Meyer's paper and infinite dimensional calculus on finitely additive canonical Hilbert space, Sept. 88.